

Impurity Non-Preserving 3-Point Correlators of BMN Operators from PP-Wave Holography II : Fermionic Excitations

Suguru DOBASHI *

*Department of Physics, Faculty of Science, Hokkaido University
Sapporo, 060-0810, Japan*

Abstract

The holographic principle in the pp-wave limit proposed in our previous works is further confirmed by studying impurity non-preserving processes which contain a fermionic BMN operator with one scalar and one fermion impurities. We show that the previously proposed duality relation between the matrix elements of the three point interaction Hamiltonian in the holographic string field theory and the OPE coefficients in super Yang-Mills theory holds to the leading order in the large μ limit. Operator mixing is required to obtain the BMN operator of definite conformal dimension which corresponds to the string state with one scalar and one fermion excitations. The mixing term plays a crucial role for our duality relation to be valid. Our results, combined with those in the previous papers, provide a positive support that our duality relation holds for the general process regardless of the kind of impurities and of whether impurities conserve or not.

*e-mail address: doxa@particle.sci.hokudai.ac.jp

1. Introduction

Holography is the key concept to the duality between string theory and gauge theory in the AdS/CFT correspondence. The holographic relation proposed by GKP/W [1] relates the degrees of freedom in the string theory defined in the bulk AdS space and those in the gauge theory defined on the conformal boundary of the AdS spacetime. Based on this proposal, a large amount of work has been done giving many positive supports for this relation [2]. In particular, the non-renormalization property of chiral primary operators has been confirmed [3]. However in the BMN limit of the AdS/CFT correspondence [4], how this holographic relation is realized has not been clear because of the lack of the clear space-time picture.

In the previous papers [5, 6, 7, 8], we gave an answer to this problem. We showed that the PP-wave background emerges as the geometry around a *tunneling null geodesic*, which starts from a point on the AdS boundary and returns to another point on the boundary. This trajectory can be obtained as a dominant path in the large angular momentum limit for the path integral which is expected to give the two point function of gauge theory operators in the GKP/W prescription[†]. The string/gauge duality in the PP-wave limit can be understood as the correspondence between the oscillation modes of strings around this trajectory and gauge theory operators sitting around its endpoints on the boundary. This picture allows a natural correspondence between an S-matrix calculation on the string theory side and an operator product expansion on the gauge theory side. As for three point functions, with the assumption that the free string basis corresponds to the BMN operators of definite conformal dimension, we proposed a duality relation between the matrix elements of the string interaction Hamiltonian and the OPE coefficients.

Our proposal in [7] has an advantage over the previous two types of duality relations, presented in [10, 11] and [12], in that it is applicable to impurity non-preserving cases. Indeed, we checked that our duality relation holds for impurity non-preserving processes which consist of BMN operators with only bosonic impurities in the previous paper [8], as well as for the impurity preserving processes [7]. To establish the duality relation for impurity non-preserving processes is important for studying the unsolved problem of string/gauge correspondence at higher genus level, since impurity non-preserving processes

[†]See also [6, 9] for more general discussions.

are known to appear as intermediate states of a string loop amplitude [13, 14, 15].

The aim of this paper is to show that our proposal also holds for the impurity non-preserving cases involving fermionic impurities. Checking our duality relation for such processes is important because our proposal was originally obtained as an extension of the duality relation for chiral primary operators to the general string excitations. It is not trivial this relation is satisfied for the non-BPS sector in general. Since we have already checked our proposal for impurity preserving sector with several different kinds of impurities, including bosons and fermions, and for impurity non-preserving sector with only bosonic impurities, what remains to be studied is impurity non-preserving processes which contain fermionic impurities. Though it is desirable to study the general processes, we restrict ourself in this paper to simple impurity non-conserving processes, where the fermionic BMN operators involved contain only one scalar and one fermionic impurities. Even for these simple examples, our duality relation holds in a very non-trivial way.

When studying the duality relation, it is important how to relate the basis between the two theories. In our prescription, the correspondent on the gauge theory side to the free string basis is a set of BMN operators of definite conformal dimension. As in the case of the singlet sector of several two impurity states [16, 17, 18, 19], the operator mixing is required for the fermionic BMN operator we are now considering so that it obtains a definite conformal dimension. We will see that this operator mixing plays a crucial role for our duality relation to hold. In addition, the phase factor of the BMN operator must be properly chosen up to $\mathcal{O}(1/J)$ in order to give the correct world sheet momentum dependence.

The result of this paper, in addition to those in the previous papers, strengthens the expectation that our duality relation is valid for general processes regardless of the kind of impurities and of whether impurities conserve or not. We believe that the successful clarification of the duality relation at the first order of the string interaction lays the foundation for the study of the duality relation at higher genus level.

The paper is organized as follows. In section 2 we briefly review our proposal for the duality relation. In section 3 we calculate two examples of the string amplitude for fermionic impurity non-preserving processes. In section 4 we first identify the gauge theory operator which corresponds to the string states with one scalar and one fermion excitations. We show by explicit perturbation calculations that for the corresponding

BMN operator to obtain a definite conformal dimension, in addition to the operator with one fermion and one scalar impurities, another operator with the same classical dimension and R-charge is required. With this operator, we compute three point functions which correspond to the string amplitudes calculated in the previous section. We will see that our duality relation holds to the leading order of the effective gauge coupling and the operator required for the diagonalization plays a crucial role. We conclude in section 5. Appendix A is devoted to the details of the perturbative calculations required for the two point function of the BMN operators with one scalar and one fermion impurities. We list the explicit forms of the prefactors in Appendix B and the asymptotic behavior of the Neumann coefficients in the large μ limit in Appendix C.

2. Review of the holographic duality relation

Our duality relation at the first order of the string interaction relates the basic ingredients which characterize the three point interaction on each theory side, the matrix element of the 3-point interaction Hamiltonian H_{123} on string theory side and the OPE coefficient C_{123} on gauge theory side. Identifying the free string basis with the set of BMN operators of definite conformal dimension, we propose that the duality relation is given by

$$C_{123} = \frac{1}{\mu(\Delta_2 + \Delta_3 - \Delta_1)} \left(f \frac{J_2 J_3}{J_1} \right)^{-\frac{\Delta_2 + \Delta_3 - \Delta_1}{2}} \Gamma \left(\frac{\Delta_2 + \Delta_3 - \Delta_1}{2} + 1 \right) H_{123}, \quad (2.1)$$

where Δ_r and J_r are the conformal dimension and the U(1) R-charge, respectively, of the gauge theory operator O_r ($r = 1, 2, 3$) which corresponds to the string state $|r, J_r\rangle$ satisfying the relation $H_2^{(r)}/\mu = \Delta_r - J_r$. Here, $H_2^{(r)}$ and μ are the light-cone free Hamiltonian and the mass parameter of the string theory, respectively, and J denotes the angular momentum around the large circle of S^5 on the string theory side. The effective gauge theory coupling is related as $\lambda' = g_{\text{YM}}^2 N / J^2 = 1/(\mu p^+ \alpha')^2$ in the PP-wave limit of AdS/CFT correspondence. The definition of f in (2.1) is given by some combination of Neumann coefficients (see the reference [7]). In what follows, we need only its asymptotic behavior in the large μ limit which is given by

$$f \frac{J_2 J_3}{J_1} \rightarrow \frac{J_1}{4\pi\mu|\alpha_{(1)}|}, \quad (2.2)$$

where $\alpha_{(r)}$ denotes $\alpha_{(r)} \equiv \alpha' p_r^+$ and we take $\alpha_{(1)} (= \alpha_{(2)} - \alpha_{(3)}) < 0, \alpha_{(2)}, \alpha_{(3)} > 0$. The relation (2.1) is expected to hold to all order of $\lambda' = 1/(\mu\alpha_{(1)})^2$, but we restrict ourself to

checking the relation to the leading order in the large μ limit in this article.

The OPE coefficient C_{123} is defined as, for scalar operators,

$$\langle O_1(x_1)O_2(x_2)O_3(x_3) \rangle = \frac{C_{123}}{|x_{12}|^{\Delta_1+\Delta_2-\Delta_3}|x_{23}|^{\Delta_2+\Delta_3-\Delta_1}|x_{31}|^{\Delta_3+\Delta_1-\Delta_2}}, \quad (2.3)$$

under the normalization of $\langle O_1 O_2 \rangle = \delta_{12}/|x_{12}|^{2\Delta}$. It is known that for the impurity preserving sector, we should take into account the operator mixing of single and double trace operators [20, 21]. However, for the impurity non-preserving process, the effect of such mixing is suppressed by $1/N$, so we need not consider this type of operator mixing as long as we are interested in the planar limit. The operator mixing we should consider later to obtain the BMN operator of definite conformal dimension is the one among single-trace operators with the same conformal dimension and R-charge.

The H_{123} is the matrix element of the 3-string interaction Hamiltonian defined as

$$H_{123} = {}_{(1)}\langle 1| {}_{(2)}\langle 2| {}_{(3)}\langle 3| \frac{\sqrt{J_1 J_2 J_3}}{N} |H_3\rangle_h, \quad (2.4)$$

under the canonical normalization of the string field theory action such as

$$S_2 = \int d\tau \left[\frac{1}{2} \langle \bar{\psi} | \partial_\tau | \psi \rangle - \frac{1}{2} (\partial_\tau \langle \bar{\psi} |) | \psi \rangle + \langle \bar{\psi} | H_2 | \psi \rangle \right], \quad (2.5)$$

$$S_3 = \frac{1}{2} \int d\tau {}_{(1)}\langle \bar{\psi} | {}_{(2)}\langle \psi | {}_{(3)}\langle \psi | \frac{\sqrt{J_1 J_2 J_3}}{N} |H_3\rangle_h + h.c., \quad (2.6)$$

$$H_2^{(r)} = \frac{1}{|\alpha^{(r)}|} \sum_{n=-\infty}^{\infty} \omega_n^{(r)} (\alpha_n^{(r)\dagger} \alpha_n^{(r)} + \beta_n^{(r)\dagger} \beta_n^{(r)}), \quad \text{with} \quad \omega_n^{(r)} = \sqrt{n^2 + (\mu \alpha_{(r)})^2}. \quad (2.7)$$

We proposed that the specific form of the string interaction Hamiltonian $|H_3\rangle_h$ is the equal weight sum of the two interaction vertexes presented so far,

$$|H_3\rangle_h = \frac{1}{2} |H_3\rangle_{SV} + \frac{1}{2} |H_3\rangle_D, \quad (2.8)$$

where $|H_3\rangle_{SV}$ is constructed in [22, 23] and further developed in [24, 25, 26, 27] as a generalization of the well-known interaction Hamiltonian in the flat space to the PP-wave background, while $|H_3\rangle_D$ is proposed in [12] and its prefactor takes the form of energy difference. These two vertexes both satisfy the SUSY algebra, as well as, the continuity condition and momentum conservation. The interaction Hamiltonian of the latter type is not adopted in the flat space case because it does not affect S-matrix calculations due to the energy conservation, but in our formalism where world-sheet time is naturally

Wick-rotated, it can give indispensable contribution. The combination (2.8) is required so that the bosonic zero-mode sector should properly reduce to the form which is obtained by taking the BMN limit of the effective action for the fluctuation fields around AdS background which corresponds to the chiral primary operators [‡]. Though in [29] it is claimed that the $|H_3\rangle_D$ part has to be modified in order to respect the $U(1)_Y$ symmetry in the supergravity sector, the modification gives no effect for the process we are now considering, as for all the cases we have confirmed so far. We will discuss the processes in the non-BPS sector for which such modification cannot be ignored in the future study [30].

For the impurity preserving processes, where $\Delta_2 + \Delta_3 - \Delta_1 = 0 + \mathcal{O}(\lambda')$, the duality relation (2.1) reduces to a simple form $\mu(\Delta_2 + \Delta_3 - \Delta_1)C_{123} = H_{123}$, the one first conjectured in [31] from a different viewpoint. We checked that this relation holds to the leading order in the large μ limit for several non-trivial processes including bosonic and fermionic impurities in the previous paper [7]. For the impurity non-preserving cases, the factor f in the duality relation plays a crucial role: due to the μ dependence of Neumann coefficients, the asymptotic behaviour of H_{123} in the large μ limit is suppressed by $\lambda' = 1/(\mu p^+ \alpha')^2$ as the difference in the number of the impurities between in- and out-states increases, while the factors $(f J_2 J_3 / J_1)^{(\Delta_1 - \Delta_2 - \Delta_3)/2}$ give a compensating contribution. Indeed, in [8], we showed that the contributions from each factor on the right hand side of (2.1) nicely combined to give the OPE coefficients C_{123} which has an appropriate μ dependence. In what follows, we confirm that this is also the case for the impurity non-preserving processes which contain fermionic BMN operators.

3. String theory calculation

We consider fermionic impurity non-preserving processes on the string theory side. In this paper we restrict ourself to simple but nontrivial processes which consist only of one boson and one fermion impurities such as

$$\text{vac} \rightarrow \alpha_m^{(2)i} \beta_{\alpha_1 \alpha_2, -m}^{(2)} + \alpha_n^{(3)j} \beta_{\beta_1 \beta_2, -n}^{(3)}, \quad (3.1)$$

$$\alpha_{n_1}^{(1)i} \alpha_{-n_1}^{(1)j} \rightarrow \alpha_{n_2}^{(2)i} \beta_{\alpha_1 \alpha_2, -n_2}^{(2)} + \alpha_{n_3}^{(3)j} \beta_{\beta_1 \beta_2, -n_3}^{(3)}. \quad (3.2)$$

[‡]In the scalar impurity sector this vertex reduces to the phenomenological interaction vertex proposed in [28], which is successfully related with the genuine OPE coefficient through the relation $\mu(\Delta_2 + \Delta_3 - \Delta_1)C_{123} = H_{123}$ in the impurity preserving process.

Even for these simple processes, we will see that the characteristic property that the interaction vertex is given by the equal weight sum of $|H_3\rangle_{SV}$ and $|H_3\rangle_D$, and the non-trivial μ dependence on the right hand side of (2.1) play important roles for our duality relation to hold. In (3.1) and (3.2), $\alpha_n^{(r)i}$ ($i = 1, 2, 3, 4$) and $\beta_{\alpha_1\alpha_2,n}^{(r)}$ denote the n -th bosonic excitation mode in a scalar direction and the n -th fermionic excitation mode, respectively, in the exponential basis of r -th string which are defined in terms of sin/cos oscillators, a_{-n}/a_n (or b_{-n}/b_n for a fermion), such as

$$\alpha_0 = a_0, \quad \alpha_n = \frac{1}{\sqrt{2}}(a_n - ia_{-n}), \quad \alpha_{-n} = \frac{1}{\sqrt{2}}(a_n + ia_{-n}), \quad (3.3)$$

$$\beta_0 = b_0, \quad \beta_n = \frac{1}{\sqrt{2}}(b_n - ib_{-n}), \quad \beta_{-n} = \frac{1}{\sqrt{2}}(b_n + ib_{-n}). \quad (3.4)$$

The $SO(8)$ spinor indexes is decomposed as $SO(8) = (SU(2) \times SU(2)) \times (SU(2) \times SU(2))$, and the subscripts of $\beta_{\alpha_1\alpha_2,n}$ denotes $(\mathbf{2}, \mathbf{1}) \times (\mathbf{2}, \mathbf{1})$ sector of the decomposition $\mathbf{8}_s = (\mathbf{2}, \mathbf{1}) \times (\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2}) \times (\mathbf{1}, \mathbf{2})$, while we denote the $(\mathbf{1}, \mathbf{2}) \times (\mathbf{1}, \mathbf{2})$ sector as $\beta_{\dot{\alpha}_1\dot{\alpha}_2,n}$.

3.1 $\text{vac} \rightarrow \alpha_m^{(2)i} \beta_{\alpha_1\alpha_2,-m}^{(2)} + \alpha_n^{(3)j} \beta_{\beta_1\beta_2,-n}^{(3)}$

The matrix element of the interaction vertex for the process (3.1) is given by

$$\langle v | \beta_{\beta_1\beta_2,-n}^{(3)} \alpha_n^{(3)j} \beta_{\alpha_1\alpha_2,-m}^{(2)} \alpha_m^{(2)i} | H_3 \rangle_h, \quad (3.5)$$

where $|H_3\rangle_h$ is the equal weight sum of the $|H_3\rangle_{SV}$ and $|H_3\rangle_D$ such as in (2.8).

Let us first calculate the contribution from $|H_3\rangle_D$ of the interaction Hamiltonian $|H_3\rangle_h$. Noting that the prefactor is given by $2 \sum_{r=1}^3 \omega_n^{(r)} / \alpha_{(r)} \sim 4\mu$ due to the form of the prefactor, $|H_3\rangle_D = (H_2^{(2)} + H_2^{(3)} - H_2^{(1)})|E\rangle$, we can easily see that the contribution from the vertex $|H_3\rangle_D$ is given by

$$\langle v | \beta_{\beta_1\beta_2,-n}^{(3)} \alpha_n^{(3)j} \beta_{\alpha_1\alpha_2,-m}^{(2)} \alpha_m^{(2)i} | H_3 \rangle_D = 4\mu \times \frac{i}{2} (Q_{nm}^{32} - Q_{mn}^{23}) \tilde{N}_{-m-n}^{23} \epsilon_{\alpha_1\beta_1} \epsilon_{\alpha_2\beta_2} \delta^{ij}. \quad (3.6)$$

Here we used the the form of the overlap vertex state such as

$$|E\rangle = |E_a\rangle |E_b\rangle, \quad (3.7)$$

$$|E_b\rangle = \exp \left[\sum_{r,s=1}^3 \sum_{m,n=1}^{\infty} \left(-\frac{i}{2} Q_{mn}^{rs} (\beta_m^{(r)\dagger\alpha_1\alpha_2} \beta_{n,\alpha_1\alpha_2}^{(s)\dagger} + \beta_m^{(r)\dagger\alpha_1\alpha_2} \beta_{-n,\alpha_1\alpha_2}^{(s)\dagger} - \beta_{-m}^{(r)\dagger\alpha_1\alpha_2} \beta_{n,\alpha_1\alpha_2}^{(s)\dagger} - \beta_{-m}^{(r)\dagger\alpha_1\alpha_2} \beta_{-n,\alpha_1\alpha_2}^{(s)\dagger}) \right) \right] |E_b^0\rangle, \quad (3.8)$$

$$|E_a\rangle = \exp \left[- \sum_{r,s=1}^3 \sum_{m,n=-\infty}^{\infty} \alpha_m^{(r)\dagger} \tilde{N}_{mn}^{rs} \alpha_n^{(s)\dagger} \right] |v\rangle, \quad (3.9)$$

where $|E_b^0\rangle$ is the overlap vertex including the fermionic zero-mode sector whose detailed form does not concern the present calculation, and the definition of Q_{mn}^{rs} is

$$Q_{mn}^{rs} = e(\alpha_{(r)}) \sqrt{\frac{|\alpha_{(s)}|}{|\alpha_{(r)}|}} \left(U_{(r)}^{1/2} C^{1/2} N^{rs} C^{-1/2} C_{(s)}^{1/2} \right)_{mn}, \quad (3.10)$$

with $U_{(r)} = C^{-1}(C_{(r)} - \mu\alpha_{(r)})$, and $C_{mn} = m\delta_{mn}$, $C_{(r)mn} = \omega_{m(r)}\delta_{mn}$. The details are referred to the reference [26]. In the large μ limit, the form of Q_{nm}^{32} and Q_{mn}^{23} are given in terms of Neumann coefficients such as

$$Q_{mn}^{23} = \frac{m}{\mu y |\alpha_{(1)}|} \tilde{N}_{mn}^{23}, \quad Q_{nm}^{32} = \frac{n}{\mu(1-y) |\alpha_{(1)}|} \tilde{N}_{nm}^{32}. \quad (3.11)$$

With this relation and the asymptotic behaviour in the large μ limit of the bosonic Neumann coefficient $\tilde{N}_{mn}^{23} = \tilde{N}_{nm}^{32}$, which is listed in Appendix C, the contribution from the interaction vertex $|H_3\rangle_D$ is given by

$$\langle v | \beta_{\beta_1\beta_2, -n}^{(3)} \alpha_n^{(3)j} \beta_{\alpha_1\alpha_2, -m}^{(2)} \alpha_m^{(2)i} | H_3 \rangle_D = -i \left(\frac{m}{y} - \frac{n}{1-y} \right) \frac{1}{8\pi^2 \mu^2 |\alpha_{(1)}|^3 y(1-y)} \epsilon_{\alpha_1\beta_1} \epsilon_{\alpha_2\beta_2} \delta^{ij}.$$

Here we have used the notation $y = -\alpha_{(2)}/\alpha_{(1)}$, $(1-y) = -\alpha_{(3)}/\alpha_{(1)}$.

As for the interaction Hamiltonian $|H_3\rangle_{SV}$, there are two types of contraction,

$$[\alpha\alpha](\beta\beta) \quad \text{or} \quad [\alpha\alpha][\beta\beta], \quad (3.12)$$

where $[XY]$ and (XY) denote the contractions of X and Y through the prefactor and overlap, respectively. For example, $[\alpha\alpha](\beta\beta)$ means $\langle v | \alpha\alpha P_{SV} | v \rangle \cdot \langle v | \beta\beta | E_b \rangle$. There are no combination from $(\beta\beta)(\alpha\alpha)$, $[\beta\beta](\alpha\alpha)$, etc. because of the form of the prefactor P_{SV} presented in Appendix B.

Using the property that, for purely bosonic external states, the interaction vertex $|H_3\rangle_{SV}$ can be rewritten as

$$\begin{aligned} |H_3\rangle_{SV} &= |H_3\rangle_D - X_{\Pi}^2 |E\rangle, \\ X_{\Pi}^2 |E\rangle &= - \sum_{r,s=1}^3 \sum_{m,n=1}^{\infty} \frac{\omega_m^{(r)}}{\alpha_{(r)}} \left(\tilde{N}_{mn}^{rs} - \tilde{N}_{m-n}^{rs} \right) \left(\alpha_m^{(r)\dagger} \alpha_n^{(s)\dagger} + \alpha_{-m}^{(r)\dagger} \alpha_{-n}^{(s)\dagger} \right) |E\rangle \\ &\quad + \sum_{r,s=1}^3 \sum_{m,n=1}^{\infty} \left(\frac{\omega_m^{(r)}}{\alpha_{(r)}} + \frac{\omega_n^{(s)}}{\alpha_{(s)}} \right) \left(\tilde{N}_{mn}^{rs} - \tilde{N}_{m-n}^{rs} \right) \alpha_m^{(r)\dagger} \alpha_{-n}^{(s)\dagger} |E\rangle, \end{aligned} \quad (3.13)$$

where X_{II} is a bosonic constituent of the prefactor whose explicit form is given in Appendix B, we can see that the contribution from $[\alpha\alpha](\beta\beta)$ is given by the product of two contributions,

$$\begin{aligned}\langle v|\alpha_m^{(2)i}\alpha_n^{(3)j}P_{SV}|E\rangle &= \langle v|\alpha_{-m}^{(2)i}\alpha_{-n}^{(3)j}(2\mu - X_{\text{II}}^2)|E\rangle \\ &= -2\mu\tilde{N}_{-m-n}^{23}\delta^{ij} + \frac{\delta^{ij}}{2}\left(\frac{\omega_m^{(2)}}{\alpha_{(2)}} + \frac{\omega_m^{(3)}}{\alpha_{(3)}}\right)(\tilde{N}_{mn}^{23} - \tilde{N}_{m-n}^{23}) \\ &= -2\mu\tilde{N}_{mn}^{23}\delta^{ij},\end{aligned}\quad (3.14)$$

$$\langle v|\beta_{\beta_1\beta_2,-n}^{(3)}\beta_{\alpha_1\alpha_2,-m}^{(2)}|E\rangle = \epsilon_{\alpha_1\beta_1}\epsilon_{\alpha_2\beta_2} \times \left(-\frac{i}{2}\right)(Q_{nm}^{32} - Q_{mn}^{23}). \quad (3.15)$$

Then, the net result of $[\alpha\alpha](\beta\beta)$ is half of the contribution from $|H_3\rangle_D$,

$$\begin{aligned}[\alpha\alpha](\beta\beta) &= i\mu(Q_{nm}^{32} - Q_{mn}^{23})\tilde{N}_{mn}^{23}\epsilon_{\alpha_1\beta_1}\epsilon_{\alpha_2\beta_2}\delta^{ij} \\ &= -i\left(\frac{m}{y} - \frac{n}{1-y}\right)\frac{1}{8\pi^2\mu^2|\alpha_{(1)}|^3y(1-y)}\epsilon_{\alpha_1\beta_1}\epsilon_{\alpha_2\beta_2}\delta^{ij}.\end{aligned}\quad (3.16)$$

As for the $[\beta\beta][\alpha\alpha]$ contribution, the only relevant term in the prefactor P_{SV} is

$$-\frac{i}{4}\tilde{K}^i\tilde{K}^jY_{ij}^2 = \frac{i}{2}X_{\text{I}}^iX_{\text{II}}^jY_{ij}^2. \quad (3.17)$$

For the definition of X_{I} , X_{II} , and Y_{ij}^2 , being bosonic and fermionic constituents of the prefactor P_{SV} , see Appendix B. Noticing that $Y_{ij}^2 = Y^{\alpha_1}_{\gamma_2}Y^{\beta_1\gamma_2}\sigma_{\alpha_1\beta_1}^{ij}$ is defined in terms of cos mode basis such as $Y^{\alpha_1\alpha_2} = \sum_{n=0}^{\infty}\bar{G}_n^{(r)}b_n^{(r)\alpha_1\alpha_2\dagger}$ and that the relation between sin/cos and exponential basis is given by (3.4), the contribution of $[\beta\beta]$ contraction is

$$\begin{aligned}\langle v|\beta_{\beta_1\beta_2,-n}^{(3)}\beta_{-m,\alpha_1\alpha_2}^{(2)}Y_{ij}^2|v\rangle &= \frac{1}{2}\langle v|b_{\beta_1\beta_2,n}^{(3)}b_{\alpha_1\alpha_2,m}^{(2)}Y^{\rho_1}_{\gamma_2}Y^{\sigma_1\gamma_2}\sigma_{\rho_1\sigma_1}^{ij}|v\rangle \\ &= -\epsilon_{\alpha_2\beta_2}(\sigma^{ij})_{\alpha_1\beta_1}\bar{G}_m^{(2)}\bar{G}_n^{(3)}.\end{aligned}\quad (3.18)$$

Here, $(\sigma^{ij})_{\alpha_1\beta_1}$ is defined as $(\sigma^{ij})_{\alpha_1\beta_1} = \epsilon_{\beta_1\gamma_1}1/2(\sigma^i\bar{\sigma}^j - \sigma^j\bar{\sigma}^i)_{\alpha_1}^{\gamma_1}$. Similarly, the bosonic contraction $[\alpha\alpha]$ is given by

$$\begin{aligned}\langle v|\alpha_m^{(2)i}\alpha_n^{(3)j}X_{\text{I}}^{[i'}X_{\text{II}}^{j']}|v\rangle &= \langle v|\alpha_m^{(2)i}\alpha_n^{(3)j} \cdot -i\sum_{r,s=1}^3\sum_{m=0,n=1}^{\infty}\bar{F}_m^{(r)}\bar{F}_n^{(s)}U_n^{(s)}a_m^{(r)\dagger[i'}a_{-n}^{(s)j']\dagger}|v\rangle \\ &= \frac{1}{2}\bar{F}_m^{(2)}\bar{F}_n^{(3)}(U_m^{(2)} - U_n^{(3)})\delta^{i[i'}\delta^{j']j}.\end{aligned}\quad (3.19)$$

Combining the two contributions, (3.18) and (3.19), and using the asymptotic form for $\bar{F}$ and $\bar{G}$ listed in Appendix C, the contribution from $[\alpha\alpha][\beta\beta]$ is evaluated as

$$[\beta\beta][\alpha\alpha] = -i\left(\frac{m}{y} - \frac{n}{1-y}\right)\epsilon_{\alpha_2\beta_2}(\sigma^{ij})_{\alpha_1\beta_1}\frac{1}{16\pi^2\mu^2|\alpha_{(1)}|^3y(1-y)}. \quad (3.20)$$

Combining all the contributions, (3.13), (3.16), and (3.20), we obtain the matrix element of the interaction vertex $|H_3\rangle_h$ for the process (3.5) in the large μ limit,

$$\begin{aligned} & \langle v | \beta_{\beta_1\beta_2,-n}^{(3)} \alpha_n^{(3)i} \beta_{\alpha_1\alpha_2,-m}^{(2)} \alpha_m^{(2)i} | H_3 \rangle_h \\ &= -i \left(\frac{m}{y} - \frac{n}{1-y} \right) \frac{1}{16\pi^2 \mu^2 |\alpha_{(1)}|^3 y(1-y)} \epsilon_{\alpha_2\beta_2} \left(\frac{3}{2} \epsilon_{\alpha_1\beta_1} \delta^{ij} + \frac{1}{2} (\sigma^{ij})_{\alpha_1\beta_1} \right). \end{aligned} \quad (3.21)$$

In order to obtain the OPE coefficient according to the duality relation (2.1), the G factor

$$\begin{aligned} G &\equiv \frac{1}{\mu(\Delta_2 + \Delta_3 - \Delta_1)} \left(f \frac{J_2 J_3}{J_1} \right)^{-\frac{\Delta_2 + \Delta_3 - \Delta_1}{2}} \Gamma \left(\frac{\Delta_2 + \Delta_3 - \Delta_1}{2} + 1 \right) \frac{\sqrt{J_1 J_2 J_3}}{N} \\ &= \frac{1}{4\mu} \left(\frac{J_1}{4\pi\mu|\alpha_{(1)}|} \right)^{-2} \Gamma(3) \frac{\sqrt{J_1 J_2 J_3}}{N} \\ &= 8\pi^2 \mu |\alpha_{(1)}|^2 J_1^{-1/2} N^{-1} \sqrt{y(1-y)}, \end{aligned} \quad (3.22)$$

must be multiplied with (3.21). Therefore, the CFT coefficient C_{123} which should be obtained from the corresponding three point correlation function on the gauge theory side is expected to be

$$C_{123} = -\frac{i}{2} \left(\frac{m}{y} - \frac{n}{1-y} \right) \frac{g_{\text{YM}} J_1^{-3/2} N^{-1/2}}{\sqrt{y(1-y)}} \epsilon_{\alpha_2\beta_2} \left(\frac{3}{2} \epsilon_{\alpha_1\beta_1} + \frac{1}{2} (\sigma^{ij})_{\alpha_1\beta_1} \right). \quad (3.23)$$

We should note that the interaction vertexes $|H_3\rangle_{SV}$ and $|H_3\rangle_D$ are both necessary to obtain this results and that the G factor properly adjust the μ -dependence to give $\mathcal{O}(g_{\text{YM}}) = \mathcal{O}(1/\mu)$ result.

3.2 $\alpha_{n_1}^{(1)i} \alpha_{-n_1}^{(1)j} \rightarrow \alpha_{n_2}^{(2)i} \beta_{\alpha_1\alpha_2,-n_2}^{(2)} + \alpha_{n_3}^{(3)j} \beta_{\beta_1\beta_2,-n_3}^{(3)}$

Next, we proceed to calculate the matrix element of the string interaction vertex for the process (3.2), which is given by

$$\langle v | \beta_{\beta_1\beta_2,-n_3}^{(3)} \alpha_{n_3}^{(3)j} \beta_{\alpha_1\alpha_2,-n_2}^{(2)} \alpha_{n_2}^{(2)i} \alpha_{n_1}^{(1)i} \alpha_{-n_1}^{(1)j} | H_3 \rangle_h. \quad (3.24)$$

We restrict ourself to the case $i \neq j$ for simplicity.

In almost the same way as the previous case, the contribution from the $|H_3\rangle_D$ is given by

$$\langle v | \beta_{\beta_1\beta_2,-n_3}^{(3)} \alpha_{n_3}^{(3)j} \beta_{\alpha_1\alpha_2,-n_2}^{(2)} \alpha_{n_2}^{(2)i} \alpha_{n_1}^{(1)i} \alpha_{-n_1}^{(1)j} | H_3 \rangle_D$$

$$\begin{aligned}
&= 2\mu \times \left(-\frac{i}{2}\right) (Q_{nm}^{32} - Q_{mn}^{23})(-\tilde{N}_{n_3, -n_1}^{31})(-\tilde{N}_{n_2 n_1}^{21}) \epsilon_{\alpha_1 \beta_1} \epsilon_{\alpha_2 \beta_2} \delta^{ij} \\
&= i \frac{1}{4\pi^2 \mu |\alpha_{(1)}|^2 y(1-y)} \epsilon_{\alpha_1 \beta_1} \epsilon_{\alpha_2 \beta_2} \delta^{ij} \left(\frac{n_2}{y} - \frac{n_3}{1-y}\right) \frac{\sin^2(\pi y n_1)}{\left(n_1 - \frac{n_2}{y}\right) \left(n_1 + \frac{n_3}{1-y}\right)}, \quad (3.25)
\end{aligned}$$

where we used the asymptotic form of the Neumann coefficients in the large μ limit presented in Appendix C.

On the other hand, because of the form of the prefactor P_{SV} , for the case $i \neq j$, the possible contribution from $|H_3\rangle_{SV}$ is that only one pair of α is contracted through the prefactor and others are through the overlap vertex:

$$\left[\alpha_{n_3}^{(3)j} \alpha_{-n_1}^{(1)j}\right] \left(\alpha_{n_2}^{(2)i} \alpha_{n_1}^{(1)i}\right) \left(\beta_{\beta_1 \beta_2, -n_3}^{(3)} \beta_{\alpha_1 \alpha_2, -n_2}^{(2)}\right) \text{ or } \left(\alpha_{n_3}^{(3)j} \alpha_{-n_1}^{(1)j}\right) \left[\alpha_{n_2}^{(2)i} \alpha_{n_1}^{(1)i}\right] \left(\beta_{\beta_1 \beta_2, -n_3}^{(3)} \beta_{\alpha_1 \alpha_2, -n_2}^{(2)}\right). \quad (3.26)$$

In particular, when $i \neq j$, the contraction through the prefactor (3.17) is not possible because of the anti-symmetric property of i and j in Y_{ij}^2 and the fact that only a pair of oscillators with the same scalar index can be contracted through the overlap vertex. This type of contraction is possible when we consider, for example, the case where the excitation of the incoming string is given by $\alpha_{n_1}^{(1)i} \alpha_{-n_1}^{(1)k} |v\rangle$ ($k \neq j$) and will play an important role for our duality relation to hold.

The amplitude of the first term in (3.26) is the product of three amplitude such as

$$\langle v | \alpha_{n_3}^{(3)j} \alpha_{-n_1}^{(1)j} P_{SV} | v \rangle \cdot \langle v | \alpha_{n_2}^{(2)i} \alpha_{n_1}^{(1)i} | E \rangle \cdot \langle v | \beta_{\beta_1 \beta_2, -n_3}^{(3)} \beta_{\alpha_1 \alpha_2, -n_2}^{(2)} | E \rangle. \quad (3.27)$$

At the first factor, the prefactor P_{SV} can be replaced with $\sum_r E_r - X_{\text{II}}$ in the the same way as in the previous subsection. But in contrast to the situation there, it gives the order $\mathcal{O}(1/\mu)$ contribution as in the impurity preserving processes:

$$\langle v | \alpha_{n_3}^{(3)j} \alpha_{-n_1}^{(1)j} \sum_r E_r | E \rangle = \left(\frac{\omega_n^{(3)}}{\alpha_{(r)}} + \frac{\omega_n^{(1)}}{\alpha_{(r)}} \right) \tilde{N}_{n_3, -n_1}^{31} = \mathcal{O}(1/\mu). \quad (3.28)$$

Since the contribution from X_{II} is of the same order and the orders of the remaining contributions are

$$\langle v | \alpha_{n_2}^{(2)i} \alpha_{n_1}^{(1)i} | E \rangle = \mathcal{O}(1), \quad \langle v | \beta_{\beta_1 \beta_2, -n_3}^{(3)} \beta_{\alpha_1 \alpha_2, -n_2}^{(2)} | E \rangle = \mathcal{O}(1/\mu^2), \quad (3.29)$$

the net result of (3.27) gives a sub-leading contribution and negligible compared to the result (3.25) in the large μ limit. The same is true for the second case of (3.26).

Therefore only the interaction vertex $|H_3\rangle_D$ gives the leading contribution such as

$$\begin{aligned} & \langle v | \beta_{\beta_1\beta_2, -n_3}^{(3)} \alpha_{n_3}^{(3)j} \beta_{\alpha_1\alpha_2, -n_2}^{(2)} \alpha_{n_2}^{(2)i} \alpha_{n_1}^{(1)i} \alpha_{-n_1}^{(1)j} | H_3 \rangle_h \\ &= i \frac{1}{8\pi^2 \mu |\alpha_{(1)}|^2 y(1-y)} \epsilon_{\alpha_1\beta_1} \epsilon_{\alpha_2\beta_2} \left(\frac{n_2}{y} - \frac{n_3}{1-y} \right) \frac{\sin^2(\pi y n_1)}{\left(n_1 - \frac{n_2}{y} \right) \left(n_1 + \frac{n_3}{1-y} \right)}. \end{aligned} \quad (3.30)$$

Taking into account the G factor for this process,

$$G = \left(\frac{J_1}{4\pi\mu|\alpha_{(1)}|} \right)^{-1} \Gamma(2) \frac{\sqrt{J_1 J_2 J_3}}{N} \frac{1}{2\mu} = 2\pi |\alpha_{(1)}| J_1^{1/2} \sqrt{y(1-y)} N^{-1}, \quad (3.31)$$

the OPE coefficient predicted from the string theory side is

$$C_{123} = i \frac{g_{\text{YM}} J_1^{-1/2} N^{-1/2}}{4\pi^2 \sqrt{y(1-y)}} \epsilon_{\alpha_1\beta_1} \epsilon_{\alpha_2\beta_2} \left(\frac{n_2}{y} - \frac{n_3}{1-y} \right) \frac{\sin^2(\pi y n_1)}{\left(n_1 - \frac{n_2}{y} \right) \left(n_1 + \frac{n_3}{1-y} \right)}. \quad (3.32)$$

4. Gauge theory calculation

Now that we have calculated two examples of the three point amplitude for the impurity non-preserving process which contain fermionic impurity operators, (3.1) and (3.2), rewritten as

$$\langle v | \beta_{\beta_1\beta_2, -n}^{(3)} \alpha_n^{(3)j} \beta_{\alpha_1\alpha_2, -m}^{(2)} \alpha_m^{(2)i} | H_3 \rangle_h, \quad (4.1)$$

$$\langle v | \beta_{\beta_1\beta_2, -n_3}^{(3)} \alpha_{n_3}^{(3)j} \beta_{\alpha_1\alpha_2, -n_2}^{(2)} \alpha_{n_2}^{(2)i} \alpha_{n_1}^{(1)i} \alpha_{-n_1}^{(1)j} | H_3 \rangle_h, \quad (4.2)$$

let us proceed to calculate the corresponding gauge theory three point correlators and derive the OPE coefficients.

For this purpose, it is important to identify properly the operator which corresponds to the string state $\alpha_n^{i\dagger} \beta_{\alpha_1\alpha_2, -n}^\dagger |v\rangle$ with the angular momentum J around S^5 . According to the prescription of inserting an impurity into the vacuum state $\text{Tr}(Z^J)$ with an appropriate phase factor, the operator with the proper U(1) R-charge and the conformal dimension satisfying the relation $H/\mu = \Delta - J$ might be given by

$$\sum_{l=0}^J \exp\left(\frac{2\pi i n}{J+2}(l+1)\right) \text{Tr}(\phi^i Z^l \lambda_{r\alpha} Z^{J-l}), \quad (4.3)$$

where the subscript r of $\lambda_{r\alpha}$ is the spinor index of the $(\mathbf{2}, \mathbf{1})_{+1/2}$ sector in the decomposition of SU(4) R-symmetry vector index A of Weyl fermion λ_α^A in the $d=4$, $\mathcal{N}=4$ SYM theory, such as

$$\mathbf{4} = (\mathbf{2}, \mathbf{1})_{+1/2} \oplus (\mathbf{1}, \mathbf{2})_{-1/2}. \quad (4.4)$$

Here the subscript $\pm 1/2$ denotes the charge of $U(1)_J$ R-symmetry subgroup we have picked up [§]. However, as we will show in what follows, in order to obtain the operator of definite conformal dimension, we must consider a linear combination of (4.3) and the operator

$$(\tau^i)_r^{\dot{r}} \text{Tr}(\theta_{\dot{r}\alpha} Z^{J+1}), \quad (4.5)$$

which is the unique possibility of the operator with the same classical dimension and $U(1)$ R-charge as the operator (4.3). Here $\theta_{\dot{r}\alpha}$ is the $(\mathbf{1}, \mathbf{2})_{-1/2}$ sector in the decomposition of (4.4) and $(\tau^i)_r^{\dot{r}}$ is the Clebsch-Gordan coefficient for a $SO(4)$ vector under $SO(4)=SU(2)\times SU(2)$ decomposition.

The operator (4.5) is expected to be obtained when the SUSY transformation and the rotation in the iZ -plane of R-symmetry act on one and the same constituent Z of the vacuum state $\text{Tr}(Z^{J+2})$, while (4.3) is obtained when these two generators act on different Z 's with an appropriate phase dependence [32, 33, 34]. Though the order of the number of the latter case is $\mathcal{O}(J)$ while the former is of $\mathcal{O}(1)$, we cannot omit this term when we identify the BMN operator which corresponds to the free string basis as in the case of the singlet sector of the operator with two bosonic or fermionic impurities studied so far [16, 19, 17, 18].

In what follows, we first determine the weight of the linear combination of (4.3) and (4.5) so that the two point function of the BMN operator takes the canonical form. Next, with the proper linear combination thus determined, we calculate the gauge theory three point functions which correspond to (4.1) and (4.2), respectively. We will see that the OPE coefficients which can be read from the three point functions exactly agree with (3.23) and (3.32).

4.1 Diagonalization and the necessity of the operator mixing

The operator on the gauge theory side which corresponds to a free string state should have a definite conformal dimension and the form of the two point function of these operators takes the canonical form determined by the conformal symmetry. The desired form of the

[§]The correspondence of the spinor indexes between string and gauge theory side is $(\alpha_1, \alpha_2) \leftrightarrow (r, \alpha)$.

two point function in the present case is

$$\begin{aligned} & \langle O_{r\alpha,m}^{J,i}(x) \bar{O}_{s\dot{\alpha},n}^{J,j}(y) \rangle \\ &= -\delta^{ij} \delta_{mn} \epsilon_{rs} \frac{1}{(x-y)^{2(J+2)}} (\sigma^\mu)_{\alpha\dot{\alpha}} \frac{(x-y)^\mu}{(x-y)^2} (1 - \lambda' n^2 \ln(x-y)^2 \Lambda^2). \end{aligned} \quad (4.6)$$

In order to obtain an operator of definite conformal dimension, we consider the linear combination such as

$$O_{r\alpha,m}^{J,i} \equiv \sum_{l=0}^J q_l \text{Tr}(\phi^i Z^l \lambda_{r\alpha} Z^{J-l}) + b(\tau^i)_r^{\dot{r}} \text{Tr}(\theta_{\dot{r}\alpha} Z^{J+1}), \quad (4.7)$$

and chose the parameter b so that the two point function,

$$\begin{aligned} \langle O_{r\alpha}^{J,i} \bar{O}_{s\dot{\alpha}}^{J,j} \rangle &= \sum_{l,l'=0}^J q_l \bar{r}_{l'} \langle \text{Tr}(\phi^i Z^l \lambda_{r\alpha} Z^{J-l}) \text{Tr}(\bar{Z}^{J-l'} \bar{\lambda}_{s\dot{\alpha}} \bar{Z}^{l'} \phi^j) \rangle \\ &+ b \langle (\tau^i)_r^{\dot{r}} \text{Tr}(\theta_{\dot{r}\alpha} Z^{J+1}) \sum_{l=0}^J \bar{r}_l \text{Tr}(\bar{Z}^{J-l} \bar{\lambda}_{s\dot{\alpha}} \bar{Z}^l \phi^j) \rangle \\ &+ \bar{b} \langle \sum_{l=0}^J q_l \text{Tr}(\bar{Z}^{J-l} \bar{\lambda}_{r\dot{\alpha}} \bar{Z}^l \phi^i) (\tau^j)_s^{\dot{s}} \text{Tr}(\bar{Z}^{J+1} \bar{\theta}_{\dot{s}\dot{\alpha}}) \rangle \\ &+ |b|^2 \langle (\tau^i)_r^{\dot{r}} \text{Tr}(\theta_{\dot{r}\alpha} Z^{J+1}) (\tau^j)_s^{\dot{s}} \text{Tr}(\bar{Z}^{J+1} \bar{\theta}_{\dot{s}\dot{\alpha}}) \rangle, \end{aligned} \quad (4.8)$$

takes the canonical form of (4.6). Here we have defined the phase factor q_l and r_l such as

$$q_l \equiv \exp\left(\frac{2\pi i m}{J+2}(l+1)\right), \quad r_l \equiv \exp\left(\frac{2\pi i n}{J+2}(l+1)\right). \quad (4.9)$$

The complex conjugate with lower indexes is defined as $\bar{O}_{r\dot{\alpha}}^i \equiv \epsilon_{rs} \bar{O}_{\dot{\alpha}}^{is} \equiv \epsilon_{rs} (O_{s\alpha}^i)^*$. Note that τ and $\bar{\tau}$ are defined as

$$(\tau^i)_{r\dot{s}} \equiv (i\sigma^i, -1), \quad (\bar{\tau}^i)^{\dot{s}r} \equiv (-i\sigma^i, -1), \quad (4.10)$$

where σ^i 's are Pauli matrices, and the complex conjugate of the matrix τ is $((\tau^i)_{r\dot{s}})^* = (\bar{\tau}^i)^{\dot{s}r}$. We will call the contributions in the first, second, third and fourth line in (4.8) as $(++)$, $(-+)$, $(+-)$, and $(--)$ sector, respectively.

In order to determine b , it is enough to focus on the log divergent part in (4.6). Let us first compute the log divergences which appear in the $(++)$ sector. The divergence comes from self-energy, $4Z$ -interaction, $2\lambda Z$ -interaction, $2\phi Z$ -interaction, and $2\phi\lambda$ -interaction, which will be explained below. The calculation is partially same as in the reference [35]

where the anomalous dimension of the operator with two scalar impurities is calculated, but here the calculation becomes a little bit complicated because we must consider additional interactions which involve fermions. We calculate here $4Z$ -interaction, etc. which have already been calculated in [35] for self-containedness. The details of the calculations are referred to Appendix A.

The two point functions of Z^a 's and $\lambda_{r\alpha}^a$'s at the one-loop level are given by

$$\langle Z^a(x) \bar{Z}^b(y) \rangle = -\frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^2 N \delta_{ab} \frac{\ln(x-y)^2 \Lambda^2}{(x-y)^2}, \quad (4.11)$$

$$\langle \lambda_{r\alpha}^a(x) \bar{\lambda}_{s\dot{\alpha}}^b(y) \rangle = - \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^2 N \delta_{ab} \epsilon_{rs} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2. \quad (4.12)$$

Hereafter, we focus only on the log divergent part and omit the other terms. The net log divergence from the self-energy in the planer limit is obtained by replacing one of the free bosonic or fermionic propagator with (4.11) or (4.12), respectively, for each possible free contraction. Then we obtain

$$C_{SE} = \sum_{l=0}^J q_l \bar{r}_l \left(-\frac{1}{2} \times (J+1) + (-1) \times 1 \right) \delta_{ij} \epsilon_{rs} \Sigma, \quad (4.13)$$

where, q , r , and Σ are defined as

$$\Sigma \equiv \frac{g_{\text{YM}}^2 N}{4\pi^2} \left(\frac{N}{2} \frac{g_{\text{YM}}^2}{4\pi^2} \right)^{N+2} \frac{1}{(x-y)^{2(J+1)}} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2. \quad (4.14)$$

The contribution from the $4Z$ -interaction is obtained by replacing the two adjacent free propagators of Z and $\bar{Z}$ with the 4-point correlator calculated at one-loop order such as

$$\langle Z^{a_1}(x) Z^{a_2}(x) \bar{Z}^{b_1}(y) \bar{Z}^{b_2}(y) \rangle = \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 (f_{pa_1b_1} f_{pa_2b_2} + f_{pa_1b_2} f_{pa_2b_1}) \frac{\ln(x-y)^2 \Lambda^2}{(x-y)^4}. \quad (4.15)$$

Apart from the phase factor, the log divergence from one possible contraction through the $4Z$ -interaction is given by

$$\begin{aligned} & \frac{1}{2} \left(\frac{N}{2} \frac{g_{\text{YM}}^2}{4\pi^2} \right)^{J-1} \frac{1}{(x-y)^{2(J-1)}} \delta_{ij} \epsilon_{rs} \sigma_{\alpha\dot{\alpha}}^\mu \partial_{x^\mu} \left(\frac{g_{\text{YM}}^2}{4\pi^2 (x-y)^2} \right) \text{Tr}(T^{a_1} T^{a_2} T^{b_2} T^{b_1}) \\ & \times \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 (f_{pa_1b_1} f_{pa_2b_2} + f_{pa_1b_2} f_{pa_2b_1}) \frac{\ln(x-y)^2 \Lambda^2}{(x-y)^4} \\ & = \frac{1}{2} \delta_{ij} \epsilon_{rs} \Sigma, \end{aligned} \quad (4.16)$$

where we have used the formula

$$f_{pa_1b_2}f_{pa_2b_1}\text{Tr}(T^{a_1}T^{a_2}T^{b_1}T^{b_2}) = N\left(\frac{N}{2}\right)^3\left(1 - \frac{1}{N^2}\right) \quad (4.17)$$

$$f_{pa_1b_1}f_{pa_2b_2}\text{Tr}(T^{a_1}T^{a_2}T^{b_1}T^{b_2}) = 0. \quad (4.18)$$

When ϕ and λ sit adjacent to each other, there are $J - 1$ ways of insertion of the $4Z$ -interaction, otherwise $J - 2$ ways. Therefore, with the phase factor appropriately taken into account, the one-loop order contribution from $4Z$ -interaction is

$$C_{4Z} = \left((q_0\bar{r}_0 + q_J\bar{r}_J)(J - 1) + \sum_{l=1}^{J-1} q_l\bar{r}_l(J - 2) \right) \times \frac{1}{2}\delta_{ij}\epsilon_{rs}\Sigma. \quad (4.19)$$

In the case of $2\lambda Z$ -interaction, the weight of the log divergence is different according to the relative position of λ and Z , since the correlator of $2\lambda Z$ -interaction is given by

$$\begin{aligned} \langle \lambda_{r\alpha}^{a_1}(x)Z^{a_2}(x)\bar{\lambda}_{s\dot{\alpha}}^{b_1}(y)\bar{Z}^{b_2}(y) \rangle &= \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 \epsilon_{rs} \left(f_{pa_1b_2}f_{pa_2b_1} + \frac{1}{2}f_{pa_1b_1}f_{pa_2b_2} \right) \\ &\times \frac{1}{(x-y)^2} \sigma_{\alpha\dot{\alpha}}^\mu \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2. \end{aligned} \quad (4.20)$$

In this case, there are 4 types of configuration of λ and Z for each possible replacement as depicted in Figure 1. The log divergence from the crossing configurations of λ and Z , which are depicted in Figure 1 with the relative phase dependence q or $\bar{r}$, is obtained by contracting (4.20) with $\text{Tr}(T^{a_1}T^{a_2}T^{b_1}T^{b_2})$, giving $1/2\delta_{ij}\epsilon_{rs}\Sigma$, while the others obtained by contracting (4.20) with $\text{Tr}(T^{a_1}T^{a_2}T^{b_2}T^{b_1})$, giving $1/4\delta_{ij}\epsilon_{rs}\Sigma$. Here, we have taken into account the free contractions of remaining operators. Then, the total contribution from $2\lambda Z$ -interaction is

$$C_{2\lambda Z} = \sum_{l=0}^{J-1} q_l\bar{r}_l \times \left(\frac{1}{4}(1 + q\bar{r}) + \frac{1}{2}(\bar{r} + q) \right) \delta_{ij}\epsilon_{rs}\Sigma, \quad (4.21)$$

where $q = \exp(2\pi im/(J + 2))$ and $r = \exp(2\pi in/(J + 2))$, respectively.

The correlator of the $2\phi Z$ -interaction is

$$\langle \phi^{ia_1}(x)Z^{a_2}(x)\phi^{jb_1}(y)\bar{Z}^{b_2}(y) \rangle = \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 \delta_{ij}f_{pa_1b_2}f_{pa_2b_1} \frac{\ln(x-y)^2\Lambda^2}{(x-y)^4}, \quad (4.22)$$

From this form of the 4-point correlator and the formula (4.18), we can easily see that only the crossing configuration of ϕ and Z gives non-zero contribution. So, we obtain, including




Figure 1: $2\lambda Z$ -interaction

the free contractions of the other operators, the log divergence from the $2\phi Z$ -interaction,

$$C_{2\phi Z} = \sum_{l=0}^{J-1} q_l \bar{r}_l \times (r + \bar{q}) \frac{1}{2} \delta_{ij} \epsilon_{rs} \Sigma. \quad (4.23)$$

The last contribution we should consider in the $(++)$ sector is the $2\phi\lambda$ -interaction, which occurs only when ϕ and λ impurities sit adjacent to each other. Since the four point correlation function is given by

$$\begin{aligned} & \langle \lambda_{r\alpha}^{a_1}(x) \phi^{ia_2}(x) \bar{\lambda}_{s\dot{\alpha}}^{b_1}(y) \phi^{ib_2}(y) \rangle \\ &= \frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 \left((-\tau^j \bar{\tau}^i)_{rs} f_{pa_1 b_2} f_{pa_2 b_1} + \frac{1}{2} (\tau^i \bar{\tau}^j)_{rs} f_{pa_1 a_2} f_{pb_1 b_2} + \delta_{ij} \epsilon_{rs} f_{pa_1 b_1} f_{pa_2 b_2} \right) \\ & \quad \times \frac{1}{(x-y)^2} \sigma_{\alpha\dot{\alpha}}^\mu \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2, \end{aligned} \quad (4.24)$$

we can easily see that the contribution from $2\phi\lambda$ -interaction is

$$\begin{aligned} C_{2\phi\lambda} &= (q_0 \bar{r}_0 + q_J \bar{r}_J) \left(\frac{1}{4} \delta_{ij} \epsilon_{rs} + \frac{1}{8} (\tau_i \tau_j)_{rs} \right) \Sigma \\ & \quad + (q_0 \bar{r}_J + q_J \bar{r}_0) \left(\frac{1}{2} \delta_{ij} \epsilon_{rs} + \frac{1}{8} (\tau_i \tau_j)_{rs} \right) \Sigma. \end{aligned} \quad (4.25)$$

Combining all the results so far, the net contribution from one-loop interactions in the $(++)$ sector of the two point function of the operators $O_{r\alpha}^{J,i}$ and $\bar{O}_{s\dot{\alpha}}^{J,i}$ is

$$\begin{aligned} & C_{SE} + C_{4Z} + C_{2\lambda Z} + C_{2\phi Z} + C_{2\phi\lambda} \\ &= \begin{cases} J \left(\frac{2\pi i n}{J} \right)^2 \delta_{ij} \epsilon_{rs} \Sigma + \frac{1}{2} (\tau^i \bar{\tau}^j)_{rs} \Sigma + \mathcal{O}(1/J^2), & (m = n) \\ + \frac{1}{2} (\tau^i \bar{\tau}^j)_{rs} \Sigma + \mathcal{O}(1/J^2), & (m \neq n) \end{cases} \end{aligned} \quad (4.26)$$

which shows that the two point function does not take the canonical form, therefore the operator (4.3) cannot be the correspondent of the $\alpha_n^i \beta_{r\alpha-n} |v\rangle$. Since the $\mathcal{O}(1)$ factor

$(\tau^i \bar{\tau}^j)_{rs}$ in (4.26) breaks the canonical form, the basic requirement for the BMN conjecture is not satisfied that the anomalous dimension to the first order of $\lambda' \equiv g_{\text{YM}}^2 N/J^2$ should match the first order perturbation expansion of the free string energy spectrum in terms of $1/\mu^2$, under the identification of $\lambda' = 1/(\mu p^+ \alpha')^2$.

The extra term in (4.26) is eliminated by choosing b properly. In the $(-+)$ sector, there appears the contribution from the 4-point correlator

$$\begin{aligned} \langle \lambda_{r\alpha}^{a_1}(x) \phi^{ia_2}(x) \bar{\theta}_{\dot{r}\dot{\alpha}}^{b_1}(y) \bar{Z}^{b_2}(y) \rangle &= \frac{1}{\sqrt{2}} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 (\tau^i)_{r\dot{r}} (f_{pa_1 b_2} f_{pa_2 b_1} + \frac{1}{2} f_{pa_1 a_2} f_{pb_1 b_2}) \\ &\times \frac{1}{(x-y)^2} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2. \end{aligned} \quad (4.27)$$

With the formula (4.17) and (4.18), we can easily see that each of 4 possible configurations of $\lambda(x)$, $\phi(x)$, $\bar{\theta}(y)$ and $\bar{Z}(y)$ gives the same contribution, $\sqrt{2}/8(\tau^i)_{r\dot{r}}\Sigma$, except for a phase factor. The same is true for the $(-+)$ sector. Then, the log divergences from the $(+-)$ and $(-+)$ sectors with appropriate phase dependences, are given by

$$b(q_0 + q_J) \frac{1}{2\sqrt{2}} (\tau^i \bar{\tau}^j)_{rs} \Sigma + \bar{b}(\bar{r}_0 + \bar{r}_J) \frac{1}{2\sqrt{2}} (\tau^i \bar{\tau}^j)_{rs} \Sigma. \quad (4.28)$$

As for the $(--)$ sector, in addition to the self-energy contribution,

$$-|b|^2 \left(-\frac{1}{2} \times (J+1) - 1 \times 1 \right) (\tau^i \bar{\tau}^j)_{rs} \Sigma, \quad (4.29)$$

where $-1/2$ comes from the self-energy of a boson propagator and -1 from fermion, and the $4Z$ -interaction contribution,

$$-|b|^2 \frac{1}{2} J (\tau^i \bar{\tau}^j)_{rs} \Sigma, \quad (4.30)$$

we have to consider the $2\theta Z$ -interaction which gives the log divergence,

$$\begin{aligned} \langle \theta_{r\alpha}^{a_1}(x) Z^{a_2}(x) \bar{\theta}_{\dot{r}\dot{\alpha}}^{b_1}(y) \bar{Z}^{b_2}(y) \rangle &= -\frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 (f_{pa_1 a_2} f_{pb_1 b_2} - f_{pa_1 b_1} f_{pa_2 b_2}) \\ &\times \frac{1}{(x-y)^2} \epsilon_{\dot{r}\dot{s}} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2. \end{aligned} \quad (4.31)$$

We have non-vanishing contribution only when θ and Z sit in the crossing position, and therefore the θZ -interaction gives the log divergence such as

$$-|b|^2 2 \times \frac{1}{4} (\tau^i \bar{\tau}^j)_{rs} \Sigma. \quad (4.32)$$

From (4.29), (4.30) and (4.32), the total contribution in the $(--)$ sector is

$$|b|^2(\tau^i \bar{\tau}^j)_{rs} \Sigma. \quad (4.33)$$

Combining all the contributions from the $(++)$, $(-+)$, $(+-)$ and $(--)$ sectors, the total log divergence is given by

$$\begin{cases} J \left(\frac{2\pi i n}{J} \right)^2 \delta_{ij} \epsilon_{rs} \Sigma + \frac{1}{2}(1 + \sqrt{2}(b + \bar{b}) + 2|b|^2)(\tau^i \bar{\tau}^j)_{rs} \Sigma + \mathcal{O}(1/J^2) & (m = n) \\ + \frac{1}{2}(1 + \sqrt{2}(b + \bar{b}) + 2|b|^2)(\tau^i \bar{\tau}^j)_{rs} \Sigma + \mathcal{O}(1/J^2) & (m \neq n) \end{cases} \quad (4.34)$$

from which we conclude that we should choose

$$b = -\frac{1}{\sqrt{2}}, \quad (4.35)$$

in order that the two point function of the operator $O_{rs}^{J,i}$ takes the canonical form.

Here we should comment on the choice of the phase factor of the operator $O_{r\alpha,m}^{J,i}$ in (4.7). In fact, we can obtain the operators of definite conformal dimension which are “diagonalized” up to $\mathcal{O}(1/J^2)$, even if we adopt the other phase dependence, for example, $\exp(2\pi i m l / (J + 1))$ or $\exp(2\pi i m l / J)$. However, according to [32, 33], we have chosen the phase factor so that the level matching condition for the world sheet momentum of the closed string is naturally realized when we regard the BMN operator $O_{rm}^{J,i}$ is obtained by acting the rotation in the R-symmetry space and the SUSY transformation on the vacuum state $\text{Tr}(Z^{J+2})$ accompanied by the phase factor q such as

$$\begin{aligned} \delta(O_1(x)O_2(x) \cdots O_n(x)) &= (\delta O_1(x))O_2(x) \cdots O_n(x) + qO_1(x)(\delta O_2(x)) \cdots O_n(x) \\ &+ \cdots + q^{n-1}O_1(x)O_2(x) \cdots (\delta O_n(x)). \end{aligned} \quad (4.36)$$

With this prescription, we have

$$\delta_2 \delta_1 \text{Tr}(Z^2) = \sum_{l=0}^{J+1} (q_1 q_2)^l \left[\sum_{l=0}^J q_2^{l+1} \text{Tr}((\delta_1 Z) Z^l (\delta_2 Z) Z^{J-l}) + \text{Tr}((\delta_2 \delta_1 Z) Z^{J+1}) \right]. \quad (4.37)$$

Then, the level matching condition is realized when we choose q_i such as $q_i = \exp(-2\pi i m_i / (J + 2))$ ($i = 1, 2$). As explained at the end of the next subsection, this choice of the phase factor is necessary for our duality condition to hold.

4.2 Three point function

Having properly identified the operator which corresponds to the string state with one scalar and one fermion excitation, $\alpha_n^i \beta_{r\alpha,-n} |v\rangle$, we proceed to calculate the three point functions which correspond to (4.1) and (4.2).




Figure 2: $(++)$ sector of three point function

$$4.2.1 \quad \text{vac} \rightarrow \alpha_{n_2}^{(2)i} \beta^{(2)} \alpha_1 \alpha_2, -n_2 + \alpha_{n_3}^{(3)j} \beta_{\beta_1 \beta_2, -n_3}^{(3)}$$

We first calculate the correspondent on the gauge theory side to the string three point function (4.1), such as

$$\langle \bar{O}_{vac}^{J_1}(x_1) O_{r\alpha, m}^{J_2, i}(x_2) O_{s\beta, n}^{J_3, j}(x_3) \rangle. \quad (4.38)$$

The normalization factor of each operator is chosen as

$$O_{vac}^{J_1} = \frac{1}{\sqrt{(J_1 + 1) N_0^{J_1 + 1}}} \text{Tr}(Z^{J_1 + 1}), \quad N_0 \equiv \frac{N}{2} \frac{g_{\text{YM}}^2}{4\pi^2}, \quad (4.39)$$

$$O_{s\alpha, m}^{J_r, i} = \frac{1}{\sqrt{2(J_r + 1) N_0^{J_r + 2}}} \left(\sum_{l=0}^{J_r} e^{\frac{2\pi i m(l+1)}{J_r + 2}} \text{Tr}(\phi^i Z^l \lambda_{s\alpha} Z^{J_r - l}) + b(\tau^i)_s^s \text{Tr}(\theta_s Z^{J_r + 1}) \right), \quad (4.40)$$

so that the two point function is normalized to $1/|x|^2$ for O_{vac}^J and takes the canonical form (4.6) for $O_{r\alpha, m}^{J, i}$. The coefficient b has been determined in the previous subsection as $b = -1/\sqrt{2}$. Similarly to the two point function, there are four sectors of the three point function to be calculated: three of them are depicted in the Figure 2, 3 and 4, which we also call $(++)$, $(+-)$ and $(--)$ sector, respectively, and the remaining one is the $(-+)$ sector, for which we have omitted the figure.

Apart from the normalization factors in (4.39) and (4.40), the $(++)$ sector of the three point function (4.38) at the leading order in terms of the genus and the gauge interaction expansions is given by

$$\frac{1}{4} \left(\frac{N}{2} \right)^{J_1 - 1} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^{J_1 + 1} \frac{(J_1 + 1) \delta_{ij}}{|x_{23}|^2 |x_{12}|^{2J_2} |x_{31}|^{2J_3}} (q_0 r_{J_3} \langle \text{Tr}(\lambda_{s\beta} \lambda_{r\alpha} \bar{Z}) \rangle + q_{J_2} r_0 \langle \text{Tr}(\lambda_{r\alpha} \lambda_{s\beta} \bar{Z}) \rangle), \quad (4.41)$$

where the three point correlator in this expression is given by

$$\begin{aligned} & \langle \text{Tr}(\bar{Z}^{a_1}(x_1)\lambda_{r\alpha}^{b_1}(x_2)\lambda_{s\beta}^{c_1}(x_3)) \rangle \\ &= i\sqrt{2}\frac{1}{g_{\text{YM}}^2}\left(\frac{g_{\text{YM}}^2}{4\pi^2}\right)^3 f_{a_1b_1c_1}\epsilon_{rs}(\sigma^\mu\bar{\sigma}^\nu)_{\alpha\beta}\int\frac{1}{(x_1-u)^2}\partial_{u^\mu}\frac{1}{(x_2-u)^2}\cdot\partial_{u^\nu}\frac{1}{(x_3-u)^2}. \end{aligned} \quad (4.42)$$

Integrating by part and using the formula (A.1), the symmetric part in terms of the spacetime indexes μ and ν in the integral can be evaluated as

$$\begin{aligned} & \int d^4u\frac{1}{(x_1-u)^2}\partial_{u^\mu}\frac{1}{(x_2-u)^2}\cdot\partial_{u^\mu}\frac{1}{(x_3-u)^2} \\ &= \frac{1}{2}\int d^4u\frac{1}{(x_1-u)^2}\left[\square\left(\frac{1}{(x_2-u)^2}\frac{1}{(x_3-u)^2}\right)\right. \\ & \quad \left.-\square\frac{1}{(x_2-u)^2}\cdot\frac{1}{(x_3-u)^2}-\frac{1}{(x_2-u)^2}\square\frac{1}{(x_3-u)^2}\right] \\ &= -2\pi^2\left(\frac{1}{|x_{12}|^2|x_{31}|^2}-\frac{1}{|x_{12}|^2|x_{23}|^2}-\frac{1}{|x_{23}|^2|x_{31}|^2}\right), \end{aligned} \quad (4.43)$$

Therefore, with the relations, $(\sigma^\mu\sigma^\nu)_{\alpha\beta} = -\eta_{\mu\nu}\epsilon_{\alpha\beta} + (\sigma^{\mu\nu})_{\alpha\beta}$, where $(\sigma^{\mu\nu})_{\alpha\beta} \equiv (\sigma^\mu\bar{\sigma}^\nu - \sigma^\nu\bar{\sigma}^\mu)_{\alpha\beta}/2$, and $f_{a_1b_1c_1}\text{Tr}(T^{a_1}T^{b_1}T^{c_1}) = iN^3/4 + \mathcal{O}(N^2)$, the $(++)$ sector of the three point function is given by

$$\begin{aligned} & \sqrt{2}\pi i\left(\frac{m}{y}-\frac{n}{1-y}\right)\left(\frac{N}{2}\frac{g_{\text{YM}}^2}{4\pi^2}\right)^{J_1+2}\delta_{ij}\epsilon_{rs}\frac{1}{|x_{12}|^{2J_2}|x_{31}|^{2J_3}|x_{23}|^2} \\ & \times\left[\epsilon_{\alpha\beta}\left(\frac{1}{|x_{12}|^2|x_{31}|^2}-\frac{1}{|x_{12}|^2|x_{23}|^2}-\frac{1}{|x_{23}|^2|x_{31}|^2}\right)\right. \\ & \quad \left.+ (\sigma^{\mu\nu})_{\alpha\beta}\int d^4u\frac{1}{(x_1-u)^2}\partial_{u^\mu}\frac{1}{(x_2-u)^2}\cdot\partial_{u^\nu}\frac{1}{(x_3-u)^2}\right]. \end{aligned} \quad (4.44)$$

The world sheet momentum dependence comes from

$$\begin{aligned} & q_1 r_{J_3+1} - q_{J_2+1} r_1 = \\ & \exp\left(\frac{2\pi i m}{J_2+2}\right)\exp\left(\frac{2\pi i n(J_3+1)}{J_3+2}\right) - \exp\left(\frac{2\pi i m(J_2+1)}{J_2+2}\right)\exp\left(\frac{2\pi i n}{J_3+2}\right) \\ & \sim 4\pi i J_1\left(\frac{m}{y}-\frac{n}{1-y}\right) \end{aligned} \quad (4.45)$$

where $y = -\alpha_{(2)}/\alpha_{(1)} = J_2/J_1$ and $1-y = -\alpha_{(3)}/\alpha_{(1)} = J_3/J_1$ and we take the leading term in the large J limit. We should note that this momentum dependence cannot be obtained if we choose the phase factor such as $\sum_{l=0}^J \exp(2\pi i n/J)\text{Tr}(\phi^i Z^l \lambda Z^{J-l})$ often adopted in the literature.




Figure 3: $(+-)$ sector of three point function




Figure 4: $(--)$ sector of three point function

As in the case of two point function, the $(+-)$ and $(-+)$ sectors are indispensable for our duality relation, while the $(--)$ sector depicted in Figure 4 gives a vanishing results since it does not contain a phase factor. The $(+-)$ sector of the three point function depicted in Figure 3 is

$$\begin{aligned} & \frac{1}{4} \left(\frac{N}{2} \right)^{J_1-1} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^{J_1+1} \frac{(J_1+1)}{|x_{12}|^{2J_2} |x_{31}|^{2J_3+2}} (\tau^j)^{\dot{s}}_s \\ & \times \left(\langle \text{Tr}(\phi^{\dot{i}}(x_2) \lambda_{r\alpha}(x_2) \theta_{\dot{s}\beta}(x_3)) \rangle + q_J \langle \text{Tr}(\lambda_{r\alpha}(x_2) \phi^{\dot{i}}(x_2) \theta_{\dot{s}\beta}(x_3)) \rangle \right). \end{aligned} \quad (4.46)$$

The perturbation calculation to the one-loop order gives

$$\begin{aligned} & \langle \text{Tr}(\phi^{a_1 i}(x_2) \lambda_{r\alpha}^{b_1}(x_2) \theta_{\dot{s}\beta}^{c_1}(x_3)) \rangle \\ & = \frac{1}{g_{\text{YM}}^2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 i f_{a_1 b_1 c_1} (\tau^i)_{r\dot{s}} (\sigma^\mu \bar{\sigma}^\nu)_{\alpha\beta} \int d^4 u \frac{1}{(x_2 - u)^2} \partial_{u^\mu} \frac{1}{(x_2 - u)^2} \cdot \partial_{u^\nu} \frac{1}{(x_3 - u)^2}. \end{aligned} \quad (4.47)$$

This time, since the antisymmetric part in terms of the indexes μ and ν can be eliminated,

the integral is transformed as

$$\begin{aligned} \frac{1}{2}(\sigma^\mu \bar{\sigma}^\nu)_{\alpha\beta} \int d^4u \partial_{u^\mu} \frac{1}{(x_2 - u)^4} \cdot \partial_{u^\nu} \frac{1}{(x_3 - u)^2} &= \frac{1}{2}\epsilon_{\alpha\beta} \int d^4u \frac{1}{(x_2 - u)^4} \square \frac{1}{(x_3 - u)^2} \\ &= -2\pi^2 \epsilon_{\alpha\beta} \frac{1}{(x_2 - x_3)^4}, \end{aligned} \quad (4.48)$$

and then the $(+-)$ sector of the three point function becomes

$$ib\pi \frac{m}{y} \left(\frac{N}{2} \frac{g_{\text{YM}}^2}{4\pi^2} \right)^{J_1+2} (\tau^i \bar{\tau}^j)_{rs} \epsilon_{\alpha\beta} \frac{1}{|x_{12}|^{2J_2} |x_{31}|^{2J_3+2} |x_{23}|^4}. \quad (4.49)$$

Similarly, the amplitude of the $(-+)$ sector is given by

$$-ib\pi \frac{n}{1-y} \left(\frac{N}{2} \frac{g_{\text{YM}}^2}{4\pi^2} \right)^{J_1+2} (\tau^i \bar{\tau}^j)_{rs} \epsilon_{\alpha\beta} \frac{1}{|x_{12}|^{2J_2+2} |x_{31}|^{2J_3} |x_{23}|^4}. \quad (4.50)$$

In the situation we are now considering, where $\epsilon = |x_2 - x_3| \ll |x_1 - x_2|$, the integral in (4.44) can be neglected because

$$(\sigma^{\mu\nu})_{\alpha\beta} \int d^4u \frac{1}{(x_1 - u)^2} \partial_{u^\mu} \frac{1}{(x_2 - u)^2} \cdot \partial_{u^\nu} \frac{1}{(x_3 - u)^2} \rightarrow 0, \quad (x_3 \rightarrow x_2). \quad (4.51)$$

Therefore, combining all the contributions, (4.44), (4.49) and (4.50), and taking into account the normalization factors in (4.39) and (4.40), the three point functions where $O_{s\beta,n}^j(x_3)$ sits in the neighbourhood of $O_{r\alpha,m}^i(x_2)$ is given by

$$\begin{aligned} &\langle \bar{O}_{vac}^{J_1}(x_1) O_{r\alpha,m}^{J_2,i}(x_2) O_{s\beta,n}^{J_3,j}(x_2 + \epsilon) \rangle \\ &= -\frac{i}{2} \left(\frac{m}{y} - \frac{n}{1-y} \right) \frac{g_{\text{YM}} J_1^{-3/2} N^{-1/2}}{\sqrt{y(1-y)}} \frac{1}{|x_{12}|^{2J_1+2} \epsilon^4} \epsilon_{\alpha\beta} \left(\left(2 + \frac{b}{\sqrt{2}} \right) \delta_{ij} \epsilon_{rs} - \frac{b}{\sqrt{2}} (\tau^{ij})_{rs} \right), \end{aligned} \quad (4.52)$$

where $(\tau^{ij})_{rs}$ is defined as $(\tau^{ij})_{rs} = \epsilon_{st} 1/2 (\tau^i \bar{\tau}^j - \tau^j \bar{\tau}^i)_r^t$. The final result (4.52) shows that the OPE coefficient which can be obtained in the limit $x_3 \rightarrow x_2$ exactly agrees with the expected result from the string theory calculation, (3.23), just when $b = -1/\sqrt{2}$. This non-trivial matching confirms that the “diagonalization” up to $\mathcal{O}(1/J)$ is crucial for our duality relation.

4.2.2 $\alpha_{n_1}^{(1)i} \alpha_{-n_1}^{(1)j} \rightarrow \alpha_{n_2}^{(2)i} \beta_{\alpha_1 \alpha_2, -n_2}^{(2)} + \alpha_{n_3}^{(3)j} \beta_{\beta_1 \beta_2, -n_3}^{(3)}$

Next, we consider the gauge theory correlation function corresponding to the process (4.2),

$$\langle \bar{O}_{n_1}^{J_1,ij}(x_1) O_{r\alpha,n_2}^{J_2,i}(x_2) O_{s\beta,n_3}^{J_3,j}(x_3) \rangle, \quad (4.53)$$




Figure 5: $(++)$ sector of three point function

at the leading order in terms of the genus and the gauge interaction expansion. Here the operator $O_{n_1}^{J_1, ij}(x_1)$ is the two-scalar impurity state defined as

$$O_{n_1}^{J_1, ij} = \frac{1}{\sqrt{(J_1 + 2)N_0^{J_1+3}}} \sum_{l_1=0}^{J_1} e^{\frac{2\pi i n_1 (l_1+1)}{J_1+2}} \text{Tr}(\phi^i Z^{l_1} \phi^j Z^{J_1-l_1}). \quad (4.54)$$

We consider the case where $i \neq j$ for simplicity as in the corresponding calculation on the string theory side. Since in this case too the $(--)$ sector does not contribute to the lowest order calculation with respect to the gauge theory interaction, it is enough to consider the $(++)$, $(+-)$, and $(-+)$ sectors.

The $(++)$ sector of this correlator at the planer level is depicted in Figure 5 and given by

$$\begin{aligned} & \frac{1}{4} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^{J_1+2} \left(\frac{N}{2} \right)^{J_1} \frac{1}{|x_{12}|^{2(J_2+1)} |x_{13}|^{2(J_3+1)}} \sum_{l_2, l_3=0}^{J_2, J_3} e^{\frac{2\pi i (l_2+1)}{J_2+2} \left(n_2 - \frac{J_2+2}{J_1+2} n_1 \right)} e^{\frac{2\pi i (l_3+1)}{J_3+2} \left(n_3 - \frac{J_3+2}{J_1+2} n_1 \right)} \\ & \times \left(e^{-\frac{2\pi i n_1 J_3}{J_1+2}} \langle \text{Tr}(\lambda_{r\alpha}(x_2) \bar{Z}(x_1) \lambda_{s\beta}(x_3)) \rangle + e^{-\frac{2\pi i n_1 (J_3+1)}{J_1+2}} \langle \text{Tr}(\bar{Z}(x_1) \lambda_{r\alpha}(x_2) \lambda_{s\beta}(x_3)) \rangle \right). \end{aligned} \quad (4.55)$$

The vacuum expectation value of $\text{Tr}(\lambda_{r\alpha}(x_2) \bar{Z}(x_1) \lambda_{s\beta}(x_3))$ in (4.55) can be evaluated in the same way as (4.42). The sum of the phase factor at the leading order of the large J limit is estimated as

$$\sum_{l_2, l_3=0}^{J_2, J_3} e^{\frac{2\pi i (l_2+1)}{J_2+2} \left(n_2 - \frac{J_2+2}{J_1+2} n_1 \right)} e^{\frac{2\pi i (l_3+1)}{J_3+2} \left(n_3 - \frac{J_3+2}{J_1+2} n_1 \right)} = -\frac{J_1}{\pi^2} e^{-2\pi i \frac{J_2+2}{J_1+2} n_1} \frac{\sin^2(\pi y n_1)}{\left(n_1 - \frac{n_2}{y} \right) \left(n_1 + \frac{n_3}{1-y} \right)}. \quad (4.56)$$




Figure 6: $(+-)$ sector of three point function

Then, the $(++)$ sector of the three point function in the situation $|x_{23}| = \epsilon \ll |x_{12}|$ is given by

$$i\sqrt{2}\frac{g_{\text{YM}}^2}{4\pi^2}\left(\frac{g_{\text{YM}}^2 N}{4\pi^2}\frac{1}{2}\right)^{J_1+3}\epsilon_{rs}\frac{1}{|x_{12}|^{2(J_1+3)}\epsilon^2}\frac{J_1}{\pi}\frac{n_1\sin^2(\pi y n_1)}{\left(n_1 - \frac{n_2}{y}\right)\left(n_1 + \frac{n_3}{1-y}\right)}. \quad (4.57)$$

On the other hand, the $(+-)$ sector in Figure 6 is

$$b\frac{1}{4}\left(\frac{g_{\text{YM}}^2}{4\pi^2}\right)^{J_1+2}\left(\frac{N}{2}\right)^{J_1}\frac{1}{|x_{12}|^{2(J_2+1)}|x_{13}|^{2(J_3+1)}}\sum_{l_2=0}^{J_2}e^{\frac{2\pi i(l_2+1)}{J_2+2}\left(n_2 - \frac{J_2+2}{J_1+2}n_1\right)}(\tau^j)_s^{\dot{s}} \quad (4.58)$$

$$\times \left(\langle \text{Tr}(\phi^j(x_1)\lambda_{r\alpha}(x_2)\theta_{\dot{s}\beta}(x_3)) \rangle + e^{-\frac{2\pi i n_1(J_3+1)}{J_1+2}} \langle \text{Tr}(\theta_{\dot{s}\beta}(x_3)\lambda_{r\alpha}(x_2)\phi^j(x_1)) \rangle \right).$$

The vacuum expectation value $\langle \text{Tr}(\phi^j(x_1)\lambda_{r\alpha}(x_2)\theta_{\dot{s}\beta}(x_3)) \rangle$ can be calculated in the same way as (4.47), and the sum of the phase factor in the large J limit is

$$\sum_{l_2=0}^{J_2}e^{\frac{2\pi i(l_2+1)}{J_2+2}\left(n_2 - \frac{J_2+2}{J_1+2}n_1\right)} = \frac{J_1}{2\pi i}\frac{1}{n_1 - \frac{n_2}{y}}(1 - e^{-2\pi i y n_1}). \quad (4.59)$$

Therefore, the result for the $(+-)$ sector is

$$ib\frac{g_{\text{YM}}^2}{4\pi^2}\left(\frac{g_{\text{YM}}^2 N}{4\pi^2}\frac{1}{2}\right)^{J_1+3}\frac{1}{|x_{12}|^{2(J_1+3)}\epsilon^2}(\tau^j\bar{\tau}^j)_{rs}\frac{J_1}{\pi}\frac{\sin^2(\pi y n_1)}{n_1 - \frac{n_2}{y}}, \quad (4.60)$$

where we do not take the sum over the superscript j . A parallel computation for the $(-+)$ sector gives

$$ib\frac{g_{\text{YM}}^2}{4\pi^2}\left(\frac{g_{\text{YM}}^2 N}{4\pi^2}\frac{1}{2}\right)^{J_1+3}\frac{1}{|x_{12}|^{2(J_1+3)}\epsilon^2}(\tau^i\bar{\tau}^i)_{rs}\frac{J_1}{\pi}\frac{\sin^2(\pi y n_1)}{n_1 + \frac{n_3}{1-y}}. \quad (4.61)$$

With the relation $(\tau^i \bar{\tau}^i)_{rs} = -\epsilon_{rs}$ (with no sum over i), and taking into account the normalization factors of BMN operators, we obtain

$$\begin{aligned}
& \langle \bar{O}_{n_1}^{J_1, ij}(x_1) O_{r\alpha, n_2}^{J_2, i}(x_2) O_{s\beta, n_3}^{J_3, j}(x_2 + \epsilon) \rangle \\
&= \frac{\sqrt{2} g_{\text{YM}} J_1^{-1/2} N^{-1/2}}{4\pi^2 \sqrt{y(1-y)}} \frac{1}{|x_{12}|^{2(J_1+3)} \epsilon^2} \epsilon_{rs} \epsilon_{\alpha\beta} \\
&\times \frac{\sin^2(\pi y n_1)}{\left(n_1 - \frac{n_2}{y}\right) \left(n_1 + \frac{n_3}{1-y}\right)} \left[2 \left(i \frac{1}{\sqrt{2}} + ib \right) n_1 - ib \left(\frac{n_2}{y} - \frac{n_3}{1-y} \right) \right]. \quad (4.62)
\end{aligned}$$

This shows that the contribution from all sectors nicely combine to match exactly the result (3.32) when $b = -1/\sqrt{2}$. As in the previous example, the non-trivial matching is realized for the operators of definite conformal dimension.

It is not difficult to consider a more general case where $O_{n_1}^{J_1, ij}$ is replaced by $O_{n_1}^{J_1, kl}$. In this case, the interaction Hamiltonian $|H_3\rangle_{SV}$ contributes to the calculation on the string theory side and gives the term proportional to $\delta^{ik} \tau_{rs}^{jl}$ or $\delta^{jl} \tau_{rs}^{ik}$ (or $(i \leftrightarrow j)$ counterparts). We can easily see that these terms also appear in the corresponding gauge theory calculation and will give the complete agreement. This gives another verification of the necessity for the both contributions from the vertexes $|H_3\rangle_D$ and $|H_3\rangle_{SV}$. We should note that when $k = l$, the operator mixing between a operator with scalar two impurities and the operator $\text{Tr}(\bar{Z} Z^{J+1})$ will also play an important role for the duality relation.

Finally, we should comment on the normalization and the phase factor of the BMN operator $O_{r\alpha, m}^{J, i}$. The normalization has been chosen so that two point function takes the form (4.6). With this choice of the normalization, our duality relation holds for the case (4.53) regardless of the choice of the phase factor, that is, whether we chose $\exp(2\pi i m/J)$, $\exp(2\pi i m/(J+1))$, or $\exp(2\pi i m/(J+2))$. On the other hand, with the normalization thus fixed, in order for our duality relation to be valid, we must chose the phase factor $\exp(2\pi i m/(J+2))$; otherwise we cannot obtain the result with the appropriate world-sheet momentum dependence and the normalization factor, as is easily seen from the calculation (4.45). Two simple examples we studied here is enough to fix both the normalization and the phase factor of the BMN operator $O_{rs, m}^{J, i}$.

5. Conclusion and Discussion

In this paper we confirm that the duality relation we proposed in [7] holds for the impurity non-conserving process which contains fermionic BMN operators. In the process of the proof, it is crucial that (a) the operator correspondent to the free string basis has the definite conformal dimensions, (b) the string interaction Hamiltonian takes the form of the equal weight sum of the two interaction vertex $|H_3\rangle_{SV}$ and $|H_3\rangle_D$ and (c) the structure of G factor in (3.22), in particular, the factor f balances the μ dependence between C_{123} and H_{123} especially in the impurity non-preserving processes.

However, though our duality relation holds in a very non-trivial way with the choice of the equal weight sum of $|H_3\rangle_{SV}$ and $|H_3\rangle_D$, we cannot conclude that this form of the string interaction vertex has proved to be completely correct. Since we cannot uniquely fix the form of the interaction vertex only from the requirement to respect the SUSY in the PP-wave background, it is not enough to check only a particular sector. In the previous paper [7] we have fixed the form of the interaction vertex so that it is consistent with the effective action for the chiral primary fields constructed in [3]. However, it has been claimed that the holographic string interaction vertex should be further modified in order to respect the $U(1)_Y$ symmetry existing in the supergravity sector [29]. They claimed that the $|H_3\rangle_D$ should be replaced as

$$|H_3\rangle_D = \sum_{r=1}^3 H_2^{(r)} |E\rangle \rightarrow \sum_{r=1}^3 H_2^{(r)} \left(1 - \frac{1}{12} Y^4\right) \left(1 - \frac{1}{12} Z^4\right) |E\rangle. \quad (5.1)$$

The new terms do not affect all the processes we have studied in previous works. The simplest example to test this modification is a impurity non-preserving process where the vacuum state splits into two BMN operators with two fermionic impurities. For this process, in contrast to the impurity preserving process where the energy difference gives $\mathcal{O}(1/\mu)$ contributions, the new term $\sum_r H_2^{(r)} Y^4 |E\rangle$ can play a significant role. In fact, it seems that for the process $\langle v | b_{11,m}^{(2)} b_{12,-m}^{(2)} b_{21,n}^{(3)} b_{22,-n}^{(3)} | H \rangle_h$ the new term cancels the contribution from the prefactor in $|H_3\rangle_{SV}$ and only $|H_3\rangle_D$ contribution survives, which is of sub-leading order in the large μ limit compared with $|H_3\rangle_{SV}$. For our duality relation to be valid, this cancellation seems to be necessary in order to obtain the correct μ dependence expected from the gauge theory calculation. In this case too, the proper identification of the BMN operator which corresponds to the free string states is crucial to check our

duality relation. The details will be reported in the future work [30].

Anyway, with the results obtained in the previous works [7, 8] that our duality relation holds for the impurity preserving process which consists of the BMN operators with scalar, vector and fermionic impurities, as well as, for the impurity non-preserving processes which consist of those with scalar and vector impurities, our duality relation itself is strongly expected to hold for the general processes regardless of the kind of impurities and of whether impurities conserve or not, as long as we choose the appropriate form of the interaction Hamiltonian in the PP-wave background and properly identify the dictionary between string and gauge theory basis.

Acknowledgements

We would like to thank T. Yoneya for valuable discussions and the collaboration at an early stage of this work.

A Feynman Graph

In this Appendix, we give the details of the calculation required to derive the log divergence of the two point function (4.8). As the regularization scheme, we adopt the differential regularization method developed in the reference [36]. With this method the anomalous dimension of the operator with two vector impurities is calculated in [33]. The formulae necessary for our calculation are

$$\frac{1}{(x-y)^4} = -\frac{1}{4}\square\frac{\ln((x-y)^2\Lambda^2)}{(x-y)^2}, \quad \square\frac{1}{(x-y)^2} = -4\pi^2\delta^{(4)}(x-y). \quad (\text{A.1})$$

With these formulae and partial integration, the two integrals needed in the calculations below to derive log divergences can be evaluated as

$$\int d^4u \frac{1}{(x-u)^4} \frac{1}{(y-u)^4} = 2\pi^2 \frac{\ln(x-y)^2\Lambda^2}{(x-y)^4}, \quad (\text{A.2})$$

$$\int d^4u \frac{1}{(x-u)^4} \frac{1}{(y-u)^2} = \pi^2 \frac{\ln(x-y)^2\Lambda^2}{(x-y)^2}. \quad (\text{A.3})$$

This regularization method is used in the calculations of the three point correlators in (4.42) and (4.47).

A.1 $\mathcal{N} = 4$ Super Yang-Mills Action

Here we present the $\mathcal{N} = 4$ Super Yang-Mills action in the 4 dimensional space-time and summarize the notation. The field content is 4 gauge fields A^μ ($\mu = 1, 2, 3, 4$), 6 real scalar fields ϕ^I ($I = 1, 2, 3, 4, 5, 6$) and 4 Weyl spinor fields λ_α^A ($A = 1, 2, 3, 4$). All these are in the adjoint representation such as

$$A^\mu = A^{a\mu} T^a, \quad \phi^I = \phi^{aI} T^a, \quad \lambda_\alpha^A T^a, \quad (\text{A.4})$$

where T^a satisfies

$$[T^a, T^b] = i f_{abc} T^c, \quad \text{Tr}[T^a, T^b] = \frac{1}{2} \delta^{ab}, \quad (\text{A.5})$$

with the structure constant f_{abc} for SU(N) symmetry, which satisfies

$$f_{apq} f_{bpq} = N \delta_{ab}, \quad f_{pqr} f_{pqr} = N(N^2 - 1). \quad (\text{A.6})$$

We pick up the U(1) subgroup of the R-symmetry such as $\text{SO}(6) = \text{U}(1) \times \text{SO}(4)$ for scalar field and $\text{SU}(4) = \text{U}(1) \times \text{SU}(2) \times \text{SU}(2)$ for spinor fields and decompose the field such as

$$\phi^I \rightarrow (\phi^i, Z, \bar{Z}), \quad \lambda_\alpha^A \rightarrow (\lambda_{r\alpha}, \theta_{\dot{r}\alpha}), \quad (\text{A.7})$$

where $Z \equiv (\phi^5 + i\phi^6)/\sqrt{2}$ which has the U(1) R-charge 1 under the rotation in the 56-plane. The indexes I, i and A are vector indexes of SO(6), SO(4) and SU(4) symmetry, respectively. The decomposed fermionic fields $\lambda_{r\alpha}$ and $\theta_{\dot{r}\alpha}$ denote the $(\mathbf{2}, \mathbf{1})_{+1/2}$ and $(\mathbf{1}, \mathbf{2})_{-1/2}$ sectors in the decomposition above, where $\pm 1/2$ is the charge under the U(1) R-symmetry we have picked up. With the decomposed field in (A.7) the $d = 4$, $\mathcal{N} = 4$ super Yang-Mills action is given by

$$\begin{aligned} g_{\text{YM}}^2 S = & -\frac{1}{4} F_{\mu\nu}^a F^{a\mu\nu} - \frac{1}{2} D_\mu \phi_i^a D^\mu \phi_i^a - \overline{D_\mu Z^a} D^\mu Z^a - \bar{\theta}^{ar} D_\mu \bar{\sigma}^\mu \theta_r^a + \bar{\lambda}_{\dot{r}}^a D_\mu \bar{\sigma}^\mu \lambda^{a\dot{r}} \\ & - \frac{1}{4} f_{pab} f_{pcd} \phi_i^a \phi_j^b \phi_i^c \phi_j^d - f_{pab} f_{pcd} \phi_i^a Z^b \phi_i^c \bar{Z}^d + \frac{1}{2} f_{pab} f_{pcd} Z^a \bar{Z}^b Z^c \bar{Z}^d \\ & - i f_{abc} \left[\theta_{\dot{r}}^a (\bar{\tau}^i)^{\dot{r}r} \lambda_r^b \phi_i^c + \bar{\lambda}^{ar} (\tau^i)_{r\dot{r}} \bar{\theta}^{b\dot{r}} \phi_i^c \right. \\ & \quad \left. - \frac{1}{\sqrt{2}} (\theta_{\dot{r}}^a \theta^{b\dot{r}} - \bar{\lambda}^{ar} \bar{\lambda}_{\dot{r}}^b) Z^c + \frac{1}{\sqrt{2}} (\bar{\theta}_{\dot{r}}^a \bar{\theta}^{b\dot{r}} - \lambda^{ar} \lambda_{\dot{r}}^b) \bar{Z}^c \right], \end{aligned} \quad (\text{A.8})$$

where $F_{\mu\nu} = \partial_{x^\mu} A_\nu - \partial_\nu A_\mu - i[A_\mu, A_\nu]$, $D_\mu Z = \partial_\mu Z - i[A_\mu, Z]$. The matrix σ_μ and τ_i are defined as

$$(\sigma^\mu)_{\alpha\dot{\alpha}} = (i\sigma^i, -1), \quad (\bar{\sigma}_\mu)^{\dot{\alpha}\alpha} = (-i\sigma^i, -1), \quad (\text{A.9})$$

$$(\tau^i)_{r\dot{r}} = (i\sigma^i, -1), \quad (\bar{\tau}_i)^{\dot{r}r} = (-i\sigma^i, -1). \quad (\text{A.10})$$

These satisfy the relation

$$(\sigma_\mu \bar{\sigma}_\nu)_\alpha^\beta + (\sigma_\nu \bar{\sigma}_\mu)_\alpha^\beta = 2\eta_{\mu\nu} \delta_\alpha^\beta, \quad (\bar{\sigma}_\mu \sigma_\nu)^{\dot{\alpha}}_{\dot{\beta}} + (\bar{\sigma}_\nu \sigma_\mu)^{\dot{\alpha}}_{\dot{\beta}} = 2\eta_{\mu\nu} \delta_{\dot{\alpha}}^{\dot{\beta}}, \quad (\text{A.11})$$

$$(\tau_i \bar{\tau}_j)_r^s + (\tau_j \bar{\tau}_i)_r^s = 2\delta_{ij} \delta_r^s, \quad (\bar{\tau}_i \tau_j)^{\dot{r}}_{\dot{s}} + (\bar{\tau}_j \tau_i)^{\dot{r}}_{\dot{s}} = 2\delta_{ij} \delta_{\dot{r}}^{\dot{s}}. \quad (\text{A.12})$$

Note that for the lowered indexes we have $(\tau_i \bar{\tau}_j)_{rs} + (\tau_j \bar{\tau}_i)_{rs} = -2\delta_{ij} \epsilon_{rs}$. The spinor indexes, as for both Lorentz and R-symmetry, are raised and lowered by the invariant tensor ϵ^{rs} and ϵ_{rs} respectively such as

$$\xi^r \equiv \epsilon^{rs} \xi_s, \quad \xi_r = \epsilon_{rs} \xi^s, \quad \epsilon^{rs} \equiv \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \epsilon_{rs} \equiv \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad (\text{A.13})$$

and we use the usual convention as

$$\xi\eta \equiv \xi^r \eta_r = \eta\xi, \quad \bar{\xi}\bar{\eta} \equiv \xi_{\dot{r}} \eta^{\dot{r}} = \bar{\eta}\bar{\xi}. \quad (\text{A.14})$$

We have omitted the Lorentz spinor indexes in (A.8) using this abbreviation.

From the SYM action (A.8), the propagators are given, in the Euclidean space, by

$$\langle \phi^{ai}(x) \phi^{bj}(y) \rangle = \delta^{ab} \delta^{ij} \frac{g_{\text{YM}}^2}{4\pi^2} \frac{1}{(x-y)^2}, \quad \langle Z^a(x) Z^b(y) \rangle = \delta^{ab} \frac{g_{\text{YM}}^2}{4\pi^2} \frac{1}{(x-y)^2}, \quad (\text{A.15})$$

$$\langle \lambda_{r\alpha}^a(x) \bar{\lambda}_{\dot{\alpha}}^{bs}(y) \rangle = -\delta^{ab} \delta_r^s \frac{g_{\text{YM}}^2}{4\pi^2} \sigma_{\alpha\dot{\alpha}}^\mu \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right), \quad (\text{A.16})$$

$$\langle \theta_{r\alpha}^a(x) \bar{\theta}_{\dot{\alpha}}^{bs}(y) \rangle = -\delta^{ab} \delta_r^s \frac{g_{\text{YM}}^2}{4\pi^2} \sigma_{\alpha\dot{\alpha}}^\mu \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right). \quad (\text{A.17})$$

A.2 Feynman diagram

We give the details of the perturbation calculations required to derive the log divergence appearing in the two point function of the operator $O_{r\alpha,m}^i$.

A.2.1 $\langle Z^a(x) \bar{Z}^b(y) \rangle$

The diagram of fermion loop self-energy is

$$\begin{aligned}
 \begin{array}{c} Z^a \\ \downarrow \\ \lambda \circlearrowleft \lambda \\ \downarrow \\ \bar{Z}^b \end{array} &= \frac{1}{2} \times 4 \times \frac{if_{apq}}{\sqrt{2}g_{\text{YM}}^2} \frac{-if_{bpq}}{\sqrt{2}g_{\text{YM}}^2} \epsilon_{rs} \epsilon^{rs} (\sigma^\mu)^\alpha_{\dot{\beta}} (\sigma^\nu)_\alpha^{\dot{\beta}} \\
 &\times \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^4 \int d^4u d^4v \frac{1}{(x-v)^2} \frac{1}{(y-u)^2} \partial_{u^\mu} \frac{1}{(u-v)^2} \cdot \partial_{u^\nu} \frac{1}{(u-v)^2}, \quad (\text{A.18})
 \end{aligned}$$

where the factor $1/2 \times 4$ comes from the perturbation expansion and the symmetric factor. With the basic formula, $(\sigma^\mu)^\alpha_{\dot{\beta}} (\sigma^\nu)_\alpha^{\dot{\beta}} = -2\eta^{\mu\nu}$, the integrand can be transformed by partial integral such that

$$\begin{aligned}
 &\int d^4u d^4v \frac{1}{(x-v)^2} \frac{1}{(y-u)^2} \partial_{u^\mu} \frac{1}{(u-v)^2} \cdot \partial_{u^\mu} \frac{1}{(u-v)^2} \\
 &= \int d^4u d^4v \frac{1}{(x-v)^2} \frac{1}{(y-u)^2} \left(\frac{1}{2} \square \frac{1}{(u-v)^4} - \square \frac{1}{(u-v)^2} \cdot \frac{1}{(u-v)^2} \right). \quad (\text{A.19})
 \end{aligned}$$

Since we focus only on the log divergence in this paper, we neglect the second term, which also should be canceled by the contribution of the same degree of divergence from other diagrams. The log divergence of the first term in (A.19) can be estimated using the formula (A.1) and (A.3). Then, the log divergence from the diagram (A.18) is given by

$$-\frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^2 N \delta_{ab} \frac{\ln(x-y)^2 \Lambda}{(x-y)^2}. \quad (\text{A.20})$$

The diagram whose intermediate states consist of θ 's also gives the same contribution.

The contribution of the gluon emission-absorption process is

$$\begin{aligned}
 \begin{array}{c} Z^a \\ \downarrow \\ \text{wavy line} \\ \downarrow \\ \bar{Z}^b \end{array} &= \frac{1}{2} \times 2 \times \frac{f_{apq}}{g_{\text{YM}}^2} \frac{f_{bpq}}{g_{\text{YM}}^2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^4 \int d^4u d^4v \frac{1}{(u-v)^2} \frac{1}{(x-u)^2} \overleftrightarrow{\partial}_{u^\mu} \frac{1}{(u-v)^2} \overleftrightarrow{\partial}_{v^\mu} \frac{1}{(y-v)^2}. \quad (\text{A.21})
 \end{aligned}$$

Integrating by part and using the formula (A.1) and (A.3), the log divergence from gluon exchange is reduced to

$$\frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^2 N \delta_{ab} \frac{\ln(x-y)^2 \Lambda^2}{(x-y)^2}. \quad (\text{A.22})$$

The log divergence which appears in the self-energy of Z is

$$= -\frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^2 N \delta_{ab} \frac{\ln(x-y)^2 \Lambda^2}{(x-y)^2}. \quad (\text{A.23})$$

A.2.2 $\langle \lambda_{r\alpha}(x) \bar{\lambda}_{s\dot{\alpha}}(y) \rangle$

The gluon emission-absorption diagram gives

$$= \frac{1}{2} \times 2 \times \epsilon_{rs} \frac{f_{apq} f_{bqp}}{g_{\text{YM}}^2 g_{\text{YM}}^2} (\sigma^\mu \bar{\sigma}^\lambda \sigma^\nu \bar{\sigma}_\lambda \sigma^\rho)_{\alpha\dot{\alpha}} \times \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^4 \int d^4 u d^4 v \frac{1}{(u-v)^2} \partial_{x^\mu} \frac{1}{(x-u)^2} \partial_{u^\nu} \frac{1}{(u-v)^2} \partial_{v^\rho} \frac{1}{(y-v)^2}. \quad (\text{A.24})$$

With the identity

$$(\sigma^\mu \bar{\sigma}^\lambda \sigma^\nu \bar{\sigma}_\lambda \sigma^\rho)_{\alpha\dot{\alpha}} = -2(\sigma^\mu \bar{\sigma}^\nu \sigma^\rho)_{\alpha\dot{\alpha}}, \quad (\text{A.25})$$

$$\sigma^\mu \bar{\sigma}^\nu \sigma^\rho = \eta^{\mu\nu} \sigma^\rho + \eta^{\nu\rho} \sigma^\mu - \eta^{\mu\rho} \sigma^\nu - i\epsilon^{\mu\nu\rho\tau} \sigma_\tau, \quad (\text{A.26})$$

and by partial integral, this integral reduces to

$$\left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^4 \epsilon_{rs} \frac{f_{apq} f_{bqp}}{g_{\text{YM}}^2 g_{\text{YM}}^2} \left[(\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \int d^4 u \frac{1}{(x-u)^2} \frac{1}{(u-v)^4} \square \frac{1}{(y-v)^2} - i\epsilon^{\mu\nu\rho\tau} (\sigma_\tau)_{\alpha\dot{\alpha}} \partial_{x^\mu} \int d^4 u \frac{1}{(x-u)^2} \partial_{u^\nu} \partial_{u^\rho} \frac{1}{(u-v)^4} \cdot \frac{1}{(y-v)^2} \right]. \quad (\text{A.27})$$

The second term vanish because of the antisymmetric property of $\epsilon^{\mu\nu\rho\tau}$, and with the formula (A.1) and (A.3), we obtain the log divergence

$$- \frac{1}{4} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^2 N \epsilon_{rs} \delta_{ab} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2. \quad (\text{A.28})$$

There are two kinds of contribution from the Yukawa interaction, where the interme-

diate states are λ, Z and θ, ϕ , respectively. The former diagram is

$$\begin{aligned}
\text{Diagram: } \begin{array}{c} \lambda_{r\alpha}^a \\ \downarrow \\ \text{Circle with } \lambda \text{ and } \bar{\lambda}_{s\dot{\alpha}}^b \end{array} &= \frac{1}{2} \times 8 \times \frac{if_{apq}}{\sqrt{2}g_{\text{YM}}^2} \frac{-if_{bpq}}{\sqrt{2}g_{\text{YM}}^2} \epsilon_{sr} (\sigma^\mu \bar{\sigma}^\nu \sigma^\rho)_{\alpha\dot{\alpha}} \\
&\times \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^4 \int d^4u d^4v \frac{1}{(u-v)^2} \frac{1}{(x-u)^2} \frac{1}{(y-v)^2} \frac{1}{(u-v)^2} \frac{1}{(y-v)^2} \\
&= -\frac{1}{4} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^2 N \epsilon_{rs} \delta_{ab} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2, \quad (\text{A.29})
\end{aligned}$$

where the integral has been performed in the same way as the gluon emission-absorption process. The latter contribution is

$$\begin{aligned}
\text{Diagram: } \begin{array}{c} \lambda_{r\alpha}^a \\ \downarrow \\ \text{Circle with } \theta \text{ and } \bar{\lambda}_{s\dot{\alpha}}^b \end{array} &= \frac{1}{2} \times 2 \times \frac{if_{apq}}{g_{\text{YM}}^2} \frac{if_{bpq}}{g_{\text{YM}}^2} (\tau^i)_{r\dot{u}} (\bar{\tau}^i)^{\dot{u}t} \epsilon_{ts} (\sigma^\mu \bar{\sigma}^\nu \sigma^\rho)_{\alpha\dot{\alpha}} \\
&\times \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^4 \int d^4u d^4v \frac{1}{(u-v)^2} \frac{1}{(x-u)^2} \frac{1}{(y-v)^2} \frac{1}{(u-v)^2} \frac{1}{(y-v)^2}. \quad (\text{A.30})
\end{aligned}$$

Since $(\tau^i)_{r\dot{u}} (\bar{\tau}^i)^{\dot{u}t} = +4\delta_r^t$, this gives the result two times of (A.29),

$$-\frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^2 N \epsilon_{rs} \delta_{ab} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2. \quad (\text{A.31})$$

The net contribution of logarithmic divergence from the fermion self-energy is

$$\begin{aligned}
\text{Diagram: } \begin{array}{c} \lambda_{r\alpha}^a \\ \downarrow \\ \text{Shaded circle} \\ \downarrow \\ \bar{\lambda}_{s\beta}^b \end{array} &= \text{Diagram: } \begin{array}{c} \downarrow \\ \text{Wavy line} \\ \downarrow \end{array} + \text{Diagram: } \begin{array}{c} \downarrow \\ \text{Circle with } \lambda \text{ and } \bar{\lambda} \end{array} + \text{Diagram: } \begin{array}{c} \downarrow \\ \text{Circle with } \theta \text{ and } \bar{\lambda} \end{array} \\
&= -\left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^2 N \epsilon_{rs} \delta_{ab} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \times \ln(x-y)^2 \Lambda^2. \quad (\text{A.32})
\end{aligned}$$

A.2.3 $\langle Z^{a_1}(x) Z^{a_2}(x) \bar{Z}^{b_1}(y) \bar{Z}^{b_2}(y) \rangle$

The amplitude for the gluon exchange diagram is given by

$$\begin{aligned}
\text{Diagram: } \begin{array}{c} Z^{a_1} \quad Z^{a_2} \\ \downarrow \quad \downarrow \\ \text{Wavy line} \\ \downarrow \quad \downarrow \\ \bar{Z}^{b_1} \quad \bar{Z}^{b_2} \end{array} &+ \text{Diagram: } \begin{array}{c} Z^{a_1} \quad Z^{a_2} \\ \downarrow \quad \downarrow \\ \text{Wavy line} \\ \downarrow \quad \downarrow \\ \bar{Z}^{b_1} \quad \bar{Z}^{b_2} \end{array} \\
&= \frac{1}{2} \times 2 \times \left(\frac{f_{pa_1b_1} f_{pa_2b_2}}{g_{\text{YM}}^2 g_{\text{YM}}^2} + \frac{f_{pa_1b_2} f_{pa_2b_1}}{g_{\text{YM}}^2 g_{\text{YM}}^2} \right) \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^5 \\
&\times \int d^4u d^4v \frac{1}{(u-v)^2} \frac{1}{(x-u)^2} \frac{1}{(y-u)^2} \frac{1}{(x-v)^2} \frac{1}{(y-v)^2} \frac{1}{(y-v)^2} \quad (\text{A.33})
\end{aligned}$$

Performing partial integral and using the formula (A.1), the integrand can be transformed as

$$\frac{1}{2}(\square_x + \square_y)H_{xyxy} + 4\pi^2 \left(\frac{2}{(x-y)^2}(Y_{xyx} + Y_{xyy}) - X_{xyxy} \right), \quad (\text{A.34})$$

where we have defined X_{abcd} and Y_{abc} as

$$H_{abcd} = \int d^4u d^4v \frac{1}{(a-u)^2} \frac{1}{(b-u)^2} \frac{1}{(u-v)^2} \frac{1}{(c-v)^2} \frac{1}{(d-v)^2}, \quad (\text{A.35})$$

$$X_{abcd} = \int d^4u \frac{1}{(a-u)^2} \frac{1}{(b-u)^2} \frac{1}{(c-u)^2} \frac{1}{(d-u)^2}, \quad (\text{A.36})$$

$$Y_{abc} = \int d^4u \frac{1}{(a-u)^2} \frac{1}{(b-u)^2} \frac{1}{(c-u)^2}. \quad (\text{A.37})$$

As long as we are concerned with the logarithmic divergence, the first term in (A.34) can be neglected and the other terms can be evaluated with the use of the formulae (A.2) and (A.3). The result is

$$\frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 (f_{pa_1b_1}f_{pa_2b_2} + f_{pa_1b_2}f_{pa_2b_1}) \frac{\ln(x-y)^2 \Lambda^2}{(x-y)^4}. \quad (\text{A.38})$$

On the other hand, the scalar 4-point interaction is easily calculated as

$$\begin{aligned} \begin{array}{c} Z^{a_1} \quad Z^{a_2} \\ \diagdown \quad \diagup \\ \diagup \quad \diagdown \\ \bar{Z}^{b_1} \quad \bar{Z}^{b_2} \end{array} &= \frac{1}{g_{\text{YM}}^2} (f_{pa_1b_1}f_{pa_2b_2} + f_{pa_1b_2}f_{pa_2b_1}) \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^4 \int d^4u \frac{1}{(x-u)^4} \frac{1}{(y-u)^4} \\ &= \frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 (f_{pa_1b_1}f_{pa_2b_2} + f_{pa_1b_2}f_{pa_2b_1}) \frac{\ln(x-y)^2 \Lambda^2}{(x-y)^4}. \end{aligned} \quad (\text{A.39})$$

Then, we obtain

$$\begin{array}{c} Z^{a_1} \quad Z^{a_2} \\ \diagdown \quad \diagup \\ \text{[Shaded Circle]} \\ \diagup \quad \diagdown \\ \bar{Z}^{b_1} \quad \bar{Z}^{b_2} \end{array} = \begin{array}{c} \downarrow \quad \downarrow \\ \text{[Wavy Line]} \\ \downarrow \quad \downarrow \end{array} + \begin{array}{c} \downarrow \quad \downarrow \\ \text{[Triangle]} \\ \downarrow \quad \downarrow \end{array} + \begin{array}{c} \downarrow \quad \downarrow \\ \text{[Cross]} \\ \downarrow \quad \downarrow \end{array} = \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 (f_{pa_1b_1}f_{pa_2b_2} + f_{pa_1b_2}f_{pa_2b_1}) \frac{\ln(x-y)^2 \Lambda^2}{(x-y)^4} \quad (\text{A.40})$$

A.2.4 $\langle \lambda_{r\alpha}^{a_1}(x) Z^{a_2}(x) \bar{\lambda}_{s\beta}^{b_1}(y) \bar{Z}^{b_2}(y) \rangle$

The log divergence which comes through the Yukawa coupling interaction is

$$\begin{array}{c} \lambda_{r\alpha}^{a_1} \quad Z^{a_2} \\ \downarrow \quad \downarrow \\ \text{[Yukawa Vertex]} \\ \downarrow \quad \downarrow \\ \bar{\lambda}_{s\alpha}^{b_1} \quad \bar{Z}^{b_2} \end{array} = \frac{1}{2} \times 8 \times \frac{if_{pa_1b_2} - if_{pa_2b_1}}{\sqrt{2}g_{\text{YM}}^2 \sqrt{2}g_{\text{YM}}^2} \epsilon_{rs} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^5 \times (\sigma^\mu \bar{\sigma}^\nu \sigma^\rho)_{\alpha\dot{\alpha}} \int d^4u d^4v \partial_{x^\mu} \frac{1}{(x-u)^2} \frac{1}{(y-u)^2} \partial_{v^\nu} \frac{1}{(u-v)^2} \partial_{v^\rho} \frac{1}{(y-v)^2} \frac{1}{(x-v)^2}. \quad (\text{A.41})$$

The integral in the second line can be evaluated as

$$\sigma_{\alpha\dot{\alpha}}^{\mu} \left(\frac{1}{2} \square_x H'_{\mu} + 2\pi^2 \partial_{x^{\mu}} X_{xxyy} \right), \quad (\text{A.42})$$

where H'_{μ} is defined as

$$\int d^4u d^4v \frac{1}{(x-u)^2} \frac{1}{(y-u)^2} \partial_{u^{\mu}} \frac{1}{(u-v)^2} \cdot \frac{1}{(x-v)^2} \frac{1}{(y-v)^2}. \quad (\text{A.43})$$

The first term in (A.42) does not give the log divergence, and the result is

$$\left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 \epsilon_{rs} f_{pa_1b_2} f_{pa_2b_1} \frac{1}{(x-y)^2} (\sigma^{\mu})_{\alpha\dot{\alpha}} \partial_{x^{\mu}} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2. \quad (\text{A.44})$$

The gluon exchange gives the contribution

$$\begin{aligned} \begin{array}{c} \lambda_{r\alpha}^{a_1} \\ \vdots \\ \text{---} \\ \bar{\lambda}_{s\dot{\alpha}}^{b_1} \end{array} \quad \begin{array}{c} Z^{a_2} \\ \downarrow \\ \text{---} \\ \bar{Z}^{b_2} \end{array} &= \frac{1}{2} \times 2 \times \frac{f_{pa_1b_1}}{g_{\text{YM}}^2} \frac{f_{pa_2b_2}}{g_{\text{YM}}^2} \epsilon_{rs} (\sigma^{\mu} \bar{\sigma}^{\nu} \sigma^{\rho})_{\alpha\dot{\alpha}} \\ &\times \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^5 \int d^4u d^4v \partial_{x^{\mu}} \frac{1}{(x-u)^2} \partial_{u^{\rho}} \frac{1}{(y-u)^2} \frac{1}{(u-v)^2} \frac{1}{(x-v)^2} \overleftrightarrow{\partial} \frac{1}{(x-v)^2} \\ &= \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^5 \frac{f_{pa_1b_1}}{g_{\text{YM}}^2} \frac{f_{pa_2b_2}}{g_{\text{YM}}^2} \epsilon_{rs} \sigma_{\alpha\dot{\alpha}}^{\mu} \left(\frac{1}{4} \square_x \partial_{x^{\mu}} H_{xxyy} + 8\pi^2 \frac{1}{(x-y)^2} \partial_{x^{\mu}} Y_{xxy} \right) \\ &= \frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 \epsilon_{rs} f_{pa_1b_1} f_{pa_2b_2} \frac{1}{(x-y)^2} (\sigma^{\mu})_{\alpha\dot{\alpha}} \partial_{x^{\mu}} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda. \end{aligned} \quad (\text{A.45})$$

Then the net result is

$$\begin{aligned} \begin{array}{c} \lambda_{r\alpha}^{a_1} \\ \searrow \\ \text{---} \\ \bar{\lambda}_{s\dot{\alpha}}^{b_1} \end{array} \quad \begin{array}{c} Z^{a_2} \\ \downarrow \\ \text{---} \\ \bar{Z}^{b_2} \end{array} &= \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} = \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 \epsilon_{rs} \left(f_{pa_1b_2} f_{pa_2b_1} + \frac{1}{2} f_{pa_1b_1} f_{pa_2b_2} \right) \\ &\times \frac{1}{(x-y)^2} \sigma_{\alpha\dot{\alpha}}^{\mu} \partial_{x^{\mu}} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2. \end{aligned} \quad (\text{A.46})$$

A.2.5 $\langle \phi^{ia_1}(x) Z^{a_2}(x) \phi^{jb_1}(y) \bar{Z}^{b_2}(y) \rangle$

The gluon exchange gives

$$\begin{aligned} \begin{array}{c} Z^{a_1} \\ \downarrow \\ \text{---} \\ \bar{Z}^{b_1} \end{array} \quad \begin{array}{c} \phi^{ia_2} \\ \downarrow \\ \text{---} \\ \phi^{jb_2} \end{array} &= \frac{1}{2} \times 4 \times \frac{f_{pa_1b_1}}{g_{\text{YM}}^2} \frac{f_{pa_2b_2}}{2g_{\text{YM}}^2} \delta_{ij} \\ &\times \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^5 \int d^4u d^4v \frac{1}{(u-v)^2} \frac{1}{(x-u)^2} \overleftrightarrow{\partial}_{u^{\mu}} \frac{1}{(y-u)^2} \frac{1}{(x-v)^2} \overleftrightarrow{\partial}_{v^{\mu}} \frac{1}{(y-v)^2} \\ &= \frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 f_{pa_1b_1} f_{pa_2b_2} \delta_{ij} \frac{\ln(x-y)^2 \Lambda^2}{(x-y)^4} \end{aligned} \quad (\text{A.47})$$

while the scalar 4-point function is

$$\begin{array}{c} Z^{a_1} \quad \phi^{ia_2} \\ \searrow \quad \swarrow \\ \bar{Z}^{b_1} \quad \phi^{jb_2} \end{array} = \frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 (f_{pa_1b_2} f_{pa_2b_1} - f_{pa_1a_2} f_{pb_1b_2}) \delta_{ij} \frac{\ln(x-y)^2 \Lambda}{(x-y)^4} \quad (\text{A.48})$$

Then, using the Jacobi identity, we obtain

$$\begin{array}{c} Z^{a_1} \quad \phi^{ia_2} \\ \searrow \quad \swarrow \\ \bar{Z}^{b_1} \quad \phi^{jb_2} \end{array} = \begin{array}{c} \downarrow \quad \downarrow \\ \text{---} \quad \text{---} \\ \downarrow \quad \downarrow \end{array} + \begin{array}{c} \searrow \quad \swarrow \\ \swarrow \quad \searrow \end{array} = \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 f_{pa_1b_2} f_{pa_2b_1} \delta_{ij} \frac{\ln(x-y)^2 \Lambda^2}{(x-y)^4}. \quad (\text{A.49})$$

A.2.6 $\langle \lambda_{r\alpha}^{a_1}(x) \phi^{ia_2}(x) \bar{\lambda}_{s\dot{\alpha}}^{b_1}(y) \phi^{ib_2}(y) \rangle$

There are two possible contraction through Yukawa interactions:

$$\begin{array}{c} \lambda_{r\alpha}^{a_1} \quad \phi^{ia_2} \\ \searrow \quad \swarrow \\ \bar{\lambda}_{s\dot{\alpha}}^{b_1} \quad \phi^{jb_2} \end{array} \begin{array}{c} \theta \\ \text{---} \end{array} = \frac{1}{2} \times 2 \times \frac{if_{pa_1b_2}}{g_{\text{YM}}^2} \frac{if_{pb_1a_2}}{g_{\text{YM}}^2} (\bar{\tau}^i)^t_s (\tau^j)_{ri} (\sigma^\mu \bar{\sigma}^\nu \sigma^\rho)_{\alpha\dot{\alpha}} \quad (\text{A.50}) \\ \times \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^5 \int d^4u d^4v \frac{1}{(x-v)^2} \frac{1}{(y-u)^2} \partial_{x^\mu} \frac{1}{(x-u)^2} \partial_{u^\nu} \frac{1}{(u-v)^2} \partial_{v^\rho} \frac{1}{(y-v)^2} \\ = - \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^5 \frac{f_{pa_1b_2}}{g_{\text{YM}}^2} \frac{f_{pa_2b_1}}{g_{\text{YM}}^2} (\tau^j \bar{\tau}^i)_{rs} (\sigma^\mu)_{\alpha\dot{\alpha}} \left(2\pi^2 \partial_{x^\mu} X_{xxyy} + \frac{1}{2} \square_x H'_\mu \right) \\ = - \frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 (\tau^j \bar{\tau}^i)_{rs} f_{pa_1b_2} f_{pa_2b_1} \frac{1}{(x-y)^2} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2,
\end{array}$$

$$\begin{array}{c} \lambda_{r\alpha}^{a_1} \quad \phi^{ia_2} \\ \searrow \quad \swarrow \\ \bar{\lambda}_{s\dot{\alpha}}^{b_1} \quad \phi^{jb_2} \end{array} \begin{array}{c} \theta \\ \text{---} \end{array} = \frac{if_{pa_1a_2}}{g_{\text{YM}}^2} \frac{if_{pb_1b_2}}{g_{\text{YM}}^2} (\bar{\tau}^j)^t_s (\tau^i)_{ri} (\sigma^\mu \bar{\sigma}^\nu \sigma^\rho)_{\alpha\dot{\alpha}} \quad (\text{A.51}) \\ \times \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^5 \int d^4u d^4v \frac{1}{(x-u)^2} \frac{1}{(y-v)^2} \partial_{x^\mu} \frac{1}{(x-u)^2} \partial_{u^\nu} \frac{1}{(u-v)^2} \partial_{v^\rho} \frac{1}{(y-v)^2} \\ = \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^5 \frac{f_{pa_1a_2}}{g_{\text{YM}}^2} \frac{f_{pb_1b_2}}{g_{\text{YM}}^2} (\tau^i \bar{\tau}^j)_{rs} (\sigma^\mu)_{\alpha\dot{\alpha}} \pi^2 \partial_{x^\mu} X_{xxyy} \\ = \frac{1}{4} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 f_{pa_1a_2} f_{pb_1b_2} (\tau^i \bar{\tau}^j)_{rs} \frac{1}{(x-y)^2} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2
\end{array}$$

The gluon exchange is the same as (A.45),

$$\begin{array}{c} \lambda_{r\alpha}^{a_1} \\ \downarrow \\ \text{---} \\ \downarrow \\ \bar{\lambda}_{s\dot{\alpha}}^{b_1} \end{array} \begin{array}{c} \phi^{ia_2} \\ \downarrow \\ \text{---} \\ \downarrow \\ \phi^{jb_2} \end{array} = \frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 f_{pa_1b_1} f_{pa_2b_2} \epsilon_{rs} \frac{1}{(x-y)^2} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2.$$

(A.52)

Then we obtain

$$\begin{array}{c} \lambda_{r\alpha}^{a_1} \\ \downarrow \\ \text{---} \\ \downarrow \\ \bar{\lambda}_{s\dot{\alpha}}^{b_1} \end{array} \begin{array}{c} \phi^{ia_2} \\ \downarrow \\ \text{---} \\ \downarrow \\ \phi^{jb_2} \end{array} = \begin{array}{c} \text{---} \\ \downarrow \\ \theta \\ \downarrow \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ \downarrow \\ \theta \\ \downarrow \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ \downarrow \\ \text{---} \\ \downarrow \\ \text{---} \end{array}$$

(A.53)

$$= \frac{1}{2} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 \left(-(\tau^j \bar{\tau}^i)_{rs} f_{pa_1b_2} f_{pa_2b_1} + \frac{1}{2} (\tau^i \bar{\tau}^j)_{rs} f_{pa_1a_2} f_{pb_1b_2} + \delta_{ij} \epsilon_{rs} f_{pa_1b_1} f_{pa_2b_2} \right) \times \frac{1}{(x-y)^2} \sigma_{\alpha\dot{\alpha}}^\mu \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2.$$

A.2.7 $\langle \lambda_{r\alpha}^{a_1}(x) \phi^{ia_2}(x) \bar{\theta}_{s\dot{\alpha}}^{b_1}(y) \bar{Z}^{b_2}(y) \rangle$

There are two kinds of the Yukawa interaction such as

$$\begin{array}{c} \lambda_{r\alpha}^{a_1} \\ \downarrow \\ \text{---} \\ \downarrow \\ \bar{\theta}_{s\dot{\alpha}}^{b_1} \end{array} \begin{array}{c} \phi^{ia_2} \\ \downarrow \\ \text{---} \\ \downarrow \\ \bar{Z}^{b_2} \end{array} = \frac{1}{2} \times 4 \times \frac{-if_{pa_1b_2}}{\sqrt{2}g_{\text{YM}}^2} \frac{-if_{pa_2b_1}}{g_{\text{YM}}^2} (\tau^i)_{r\dot{r}} (\sigma^\mu \bar{\sigma}^\nu \sigma^\rho)_{\alpha\dot{\alpha}}$$

(A.54)

$$\times \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^5 \int d^4u d^4v \frac{1}{(x-v)^2} \frac{1}{(y-u)^2} \partial_{x^\mu} \frac{1}{(x-u)^2} \partial_{v^\nu} \frac{1}{(u-v)^2} \partial_{y^\rho} \frac{1}{(y-v)^2}$$

$$= \frac{1}{\sqrt{2}} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 (\tau^i)_{r\dot{r}} f_{pa_1b_2} f_{pa_2b_1} \frac{1}{(x-y)^2} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2$$


$$\begin{array}{c} \lambda_{r\alpha}^{a_1} \\ \downarrow \\ \text{---} \\ \downarrow \\ \bar{\theta}_{s\dot{\alpha}}^{b_1} \end{array} \begin{array}{c} \phi^{ia_2} \\ \downarrow \\ \text{---} \\ \downarrow \\ \bar{Z}^{b_2} \end{array} = \frac{1}{2} \times 4 \times (\tau^i)_{r\dot{r}} \frac{-if_{pa_1a_2}}{g_{\text{YM}}^2} \frac{if_{pb_1b_2}}{\sqrt{2}g_{\text{YM}}^2} (\sigma^\mu \bar{\sigma}^\nu \sigma^\rho)_{\alpha\dot{\alpha}}$$

(A.55)

$$\times \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^5 \int d^4u d^4v \frac{1}{(x-u)^2} \frac{1}{(y-v)^2} \partial_{x^\mu} \frac{1}{(x-u)^2} \partial_{u^\nu} \frac{1}{(u-v)^2} \partial_{y^\rho} \frac{1}{(y-v)^2}$$

$$= \frac{1}{2\sqrt{2}} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 (\tau^i)_{r\dot{r}} f_{pa_1a_2} f_{pb_1b_2} \frac{1}{(x-y)^2} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \ln(x-y)^2 \Lambda^2$$

Then the sum of the above contribution gives the result (4.27):




$$(A.56)$$

$$= \frac{1}{\sqrt{2}} \left(\frac{g_{\text{YM}}^2}{4\pi^2} \right)^3 (\tau^i)_{r\dot{r}} (f_{pa_1b_2} f_{pa_2b_1} + \frac{1}{2} f_{pa_1a_2} f_{pb_1b_2}) \frac{1}{(x-y)^2} (\sigma^\mu)_{\alpha\dot{\alpha}} \partial_{x^\mu} \left(\frac{1}{(x-y)^2} \right) \times \ln(x-y)^2 \Lambda^2. \quad (A.57)$$


A.2.8 $\langle \theta_{\dot{r}\alpha}^{a_1}(x) Z^{a_2}(x) \bar{\theta}_{\dot{s}\dot{\alpha}}^{b_1}(y) \bar{Z}^{b_2}(y) \rangle$

The contribution from the Yukawa coupling is




$$(A.58)$$

The gluon exchange is the same as (A.45),



$$(A.59)$$

Therefore we obtain



$$(A.60)$$

B Prefactor

The form of the prefactor of the interaction vertex $|H_3\rangle_{SV}$ is[¶]

$$|H_3\rangle_{SV} = P_{SV}|E\rangle, \quad (B.1)$$

$$P_{SV} = \frac{1}{2} \left[\left(K^i \tilde{K}^j + \mu \delta^{ij} \right) V_{ij} - \left(K^\mu \tilde{K}^\nu + \mu \delta^{\mu\nu} \right) V_{\mu\nu} \right. \\ \left. - K^{\dot{\alpha}_1 \alpha_1} \tilde{K}^{\dot{\alpha}_2 \alpha_2} S_{\alpha_1 \alpha_2}(Y) S_{\dot{\alpha}_1 \dot{\alpha}_2}^*(Z) - \tilde{K}^{\dot{\alpha}_1 \alpha_1} K^{\dot{\alpha}_2 \alpha_2} S_{\alpha_1 \alpha_2}^*(Y) S_{\dot{\alpha}_1 \dot{\alpha}_2}(Z) \right] \quad (B.2)$$

where K^I , $\tilde{K}^I$ ($I = i/\mu$ denotes a scalar/vector direction) and $Y^{\alpha_1 \alpha_2}$, $Z^{\dot{\alpha}_1 \dot{\alpha}_2}$ are bosonic and fermionic constituents of the prefactor defined as

$$K^J = X_I^J + X_{II}^J, \quad \tilde{K}^J = X_I^J - X_{II}^J, \quad (B.3)$$

$$X_I = \sum_{r=1}^3 \sum_{n=0}^{\infty} \bar{F}_n^{(r)} a_n^{(r)\dagger}, \quad X_{II} = \sum_{r=1}^3 \sum_{n=1}^{\infty} i U_{n(r)} \bar{F}_n^{(r)} a_{-n}^{(r)\dagger}, \quad (B.4)$$

$$Y^{\alpha_1 \alpha_2} = \sum_{r=1}^3 \sum_{n=0}^{\infty} \bar{G}_n^{(r)} b_n^{(r) \alpha_1 \alpha_2 \dagger}, \quad Z^{\dot{\alpha}_1 \dot{\alpha}_2} = \sum_{r=1}^3 \sum_{n=0}^{\infty} \bar{G}_n^{(r)} b_{-n}^{(r) \dot{\alpha}_1 \dot{\alpha}_2 \dagger}, \quad (B.5)$$

with

$$\bar{F}_n^{(r)} = \sqrt{-\frac{\alpha'}{\alpha}} (1 - 4\mu\alpha K)^{1/2} F_n^{(r)}, \quad \bar{G}_n^{(r)} = \sqrt{-\frac{\alpha'}{\alpha}} (1 - 4\mu\alpha K)^{1/2} G_n^{(r)}, \quad (B.6)$$

$$F^{(2)} = -\sqrt{\frac{2}{\alpha'}} \sqrt{\mu\alpha(2)} \alpha_{(3)}, \quad F^{(3)} = \sqrt{\frac{2}{\alpha'}} \sqrt{\mu\alpha(3)} \alpha_{(2)}, \quad F^{(1)} = 0, \quad (B.7)$$

$$F_n^{(r)} = -\frac{\alpha}{\sqrt{\alpha'}} \frac{1}{1 - 4\mu\alpha K} \frac{1}{\alpha_{(r)}} (U_{(r)}^{-1} C_{(r)}^{1/2} C N^r)_n \quad (n > 0), \quad (B.8)$$

$$G^{(2)} = -\sqrt{\frac{1}{\alpha'}} \sqrt{\alpha_{(2)}} \alpha_{(3)}, \quad G^{(3)} = \sqrt{\frac{1}{\alpha'}} \sqrt{\alpha_{(3)}} \alpha_{(2)}, \quad G^{(1)} = 0, \quad (B.9)$$

$$G_n^{(r)} = -\frac{\alpha}{\sqrt{\alpha'}} \frac{1}{1 - 4\mu\alpha K} \frac{e(\alpha_{(r)})}{\sqrt{|\alpha_{(r)}|}} (U_{(r)}^{-1/2} C_{(r)}^{1/2} C^{1/2} N^r)_n \quad (n > 0). \quad (B.10)$$

[¶]This definition differs from the one in (3.28) of [26] by a factor $-2\alpha/\alpha'$. Note also the difference of the total factor of K^i between here and there.

The other quantities in the prefactor is defined as

$$V_{ij} \equiv \delta_{ij} \left[1 + \frac{1}{12}(Y^4 + Z^4) + \frac{1}{144}Y^4 Z^4 \right] - \frac{i}{2} \left[Y_{ij}^2 (1 + \frac{1}{12}Z^4) - Z_{ij}^2 (1 + \frac{1}{12}Y^4) \right] + \frac{1}{4}(Y^2 Z^2)_{ij}, \quad (\text{B.11})$$

$$V_{\mu\nu} \equiv \delta_{\mu\nu} \left[1 - \frac{1}{12}(Y^4 + Z^4) + \frac{1}{144}Y^4 Z^4 \right] - \frac{i}{2} \left[Y_{\mu\nu}^2 (1 - \frac{1}{12}Z^4) - Z_{\mu\nu}^2 (1 - \frac{1}{12}Y^4) \right] + \frac{1}{4}(Y^2 Z^2)_{\mu\nu}, \quad (\text{B.12})$$

$$S(Y) \equiv Y + \frac{i}{3}Y^3, \quad (\text{B.13})$$

with

$$K^{\dot{\alpha}_r \beta_r} \equiv K^i \sigma_i^{\dot{\alpha}_r \beta_r}, \quad \tilde{K}^{\dot{\alpha}_r \beta_r} \equiv K^i \sigma_i^{\dot{\alpha}_r \beta_r}, \quad (r = 1, 2) \quad (\text{B.14})$$

$$Y_{\alpha_1 \beta_1}^2 \equiv Y_{\alpha_1 \alpha_2} Y_{\beta_1}^{\alpha_2}, \quad Y_{\alpha_2 \beta_2}^2 \equiv Y_{\alpha_1 \alpha_2} Y_{\beta_2}^{\alpha_1}, \quad (\text{B.15})$$

$$Y_{\alpha_1 \beta_2}^3 \equiv Y_{\alpha_1 \beta_1}^2 Y_{\beta_2}^{\beta_1}, \quad Y^4 \equiv Y_{\alpha_1 \beta_1}^2 Y^{2\alpha_1 \beta_1}, \quad (\text{B.16})$$

$$Y^{2ij} \equiv Y^{2\alpha_1 \beta_1} \sigma_{\alpha_1 \beta_1}^{ij}, \quad Z^{2ij} \equiv Z^{2\dot{\alpha}_1 \dot{\beta}_1} \sigma_{\dot{\alpha}_1 \dot{\beta}_1}^{ij}, \quad (Y^2 Z^2)^{ij} \equiv Y^{2k(i} Z^{2j)k}, \quad (\text{B.17})$$

We refer the reader to the ref.[26] for details.

On the other hand, the interaction vertex presented in [12], which is of the form

$$|H_3\rangle_D = \sum_{r=1}^3 \sum_{m=-\infty}^{\infty} \frac{\omega_m^{(r)}}{\alpha^{(r)}} \left(\sum_{I=1}^8 a_m^{(r)I\dagger} a_m^{(r)I} + \sum_{a=1}^8 b_m^{(r)a\dagger} b_m^{(r)a} \right) |E\rangle \quad (\text{B.18})$$

can be written as

$$|H_3\rangle_D = P_D |E\rangle, \quad (\text{B.19})$$

$$P_D = \frac{1}{4} \left(K^2 + \tilde{K}^2 \right) - Y^{\alpha_1 \alpha_2} \tilde{Y}_{\alpha_1 \alpha_2} - Z^{\dot{\alpha}_1 \dot{\alpha}_2} \tilde{Z}_{\dot{\alpha}_1 \dot{\alpha}_2}, \quad (\text{B.20})$$

where

$$\tilde{Y}^{\alpha_1 \alpha_2} = \sum_{r=1}^3 \frac{n}{\alpha^{(r)}} G_n^{(r)} b_n^{(r)\alpha_1 \alpha_2 \dagger}, \quad \tilde{Z}^{\dot{\alpha}_1 \dot{\alpha}_2} = \sum_{r=1}^3 \frac{n}{\alpha^{(r)}} G_n^{(r)} b_{-n}^{(r)\alpha_1 \alpha_2 \dagger}. \quad (\text{B.21})$$

C Large μ behavior

The large μ behavior of $\tilde{N}_{mn}^{rs}$ is given [37], for $(m, n) \neq (0, 0)$, by

$$\tilde{N}_{mn}^{22} = \frac{(-1)^{m+n}}{4\pi\mu|\alpha_{(1)}|y}, \quad \tilde{N}_{mn}^{23} = \frac{(-1)^{m+1}}{4\pi\mu|\alpha_{(1)}|\sqrt{y(1-y)}} \quad (\text{C.1})$$

$$\tilde{N}_{mn}^{33} = \frac{1}{4\pi\mu|\alpha_{(1)}|(1-y)}, \quad \tilde{N}_{mn}^{11} = \frac{(-1)^{m+n+1} \sin(\pi my) \sin(\pi ny)}{\pi\mu|\alpha_{(1)}|} \quad (\text{C.2})$$

$$\tilde{N}_{mn}^{21} = \frac{(-1)^{m+n+1} \sin(\pi ny)}{\pi\sqrt{y}(n-m/y)}, \quad \tilde{N}_{mn}^{31} = \frac{(-1)^n \sin(\pi ny)}{\pi\sqrt{1-y}(n-m/(1-y))} \quad (\text{C.3})$$

and, for $m = n = 0$, by

$$\tilde{N}_{00}^{11} = 0, \quad \tilde{N}_{00}^{12} = -\sqrt{y}, \quad \tilde{N}_{00}^{13} = -\sqrt{1-y}, \quad (\text{C.4})$$

$$\tilde{N}_{00}^{23} = -\frac{1}{4\pi\mu|\alpha_{(1)}|\sqrt{y(1-y)}}, \quad \tilde{N}_{00}^{22} = \frac{1}{4\pi\mu|\alpha_{(1)}|y}, \quad \tilde{N}_{00}^{33} = \frac{1}{4\pi\mu|\alpha_{(1)}|(1-y)} \quad (\text{C.5})$$

When we compute string amplitudes for fermions, the large μ behavior of $\bar{F}_n^{(r)}$, $\bar{G}_n^{(r)}$, $U_m^{(r)}$ and $1 - 4\mu\alpha K$ are also useful^{||}:

$$\bar{F}_n^{(2)} = \frac{(-1)^{n+1}\sqrt{\mu}}{\sqrt{\pi\mu|\alpha_{(1)}|y}}, \quad \bar{F}_n^{(3)} = \frac{\sqrt{\mu}}{\sqrt{\pi\mu|\alpha_{(1)}|(1-y)}}, \quad \bar{F}_n^{(1)} = \frac{(-1)^{n+1}\sqrt{\mu}n \sin(\pi ny)}{\sqrt{\pi(\mu|\alpha_{(1)}|)^3}} \quad (\text{C.6})$$

$$\bar{F}_0^{(2)} = -\frac{\sqrt{\mu}}{\sqrt{2\pi\mu|\alpha_{(1)}|y}}, \quad \bar{F}_0^{(3)} = \frac{\sqrt{\mu}}{\sqrt{2\pi\mu|\alpha_{(1)}|(1-y)}}, \quad \bar{F}_0^{(1)} = 0 \quad (\text{C.7})$$

$$\bar{G}_n^{(2)} = \frac{(-1)^{n+1}}{\sqrt{2\pi\mu|\alpha_{(1)}|y}}, \quad \bar{G}_n^{(3)} = \frac{1}{\sqrt{2\pi\mu|\alpha_{(1)}|(1-y)}}, \quad \bar{G}_n^{(1)} = \frac{(-1)^{n+1}\sqrt{2} \sin(\pi ny)}{\sqrt{\pi\mu|\alpha_{(1)}|}} \quad (\text{C.8})$$

$$\bar{G}_0^{(2)} = -\frac{1}{\sqrt{4\pi\mu|\alpha_{(1)}|y}}, \quad \bar{G}_0^{(3)} = \frac{1}{\sqrt{4\pi\mu|\alpha_{(1)}|(1-y)}}, \quad \bar{G}_0^{(1)} = 0 \quad (\text{C.9})$$

$$U_n^{(2)} = \frac{n}{2\mu|\alpha_{(1)}|y}, \quad U_n^{(3)} = \frac{n}{2\mu|\alpha_{(1)}|(1-y)}, \quad U_n^{(1)} = \frac{2\mu|\alpha_{(1)}|}{n}, \quad (\text{C.10})$$

$$1 - 4\mu\alpha K = \frac{1}{4\pi\mu|\alpha_{(1)}|y(1-y)}, \quad (\text{C.11})$$

References

- [1] S.S. Gubser, I.R. Klebanov, and A.M. Polyakov, *Gauge theory correlators from noncritical string theory*, Phys.Lett.B428(1998)105, hep-th/9802109,
- E. Witten, *Anti-de Sitter space and holography*, Adv.Theor.Math.Phys.2(1998)253, hep-th/9802150

^{||}Note that the definition of the Neumann vector N_n^r here, with which $F_n^{(r)}$ is defined, differs by $C_{(r)}^{-1/2}U_r$ from that of the ref.[37].

- [2] O. Aharony, S.S. Gubser, J.M. Maldacena, H. Ooguri, and Y. Oz, *Large N Field Theories, String Theory and Gravity*, Phys.Rept.323(2000)183, hep-th/9905111
E. D'Hoker and D.Z. Freedman, *Supersymmetric gauge theories and the AdS/CFT correspondence*, TASI 2001 Lecture Notes, hep-th/0201253
- [3] S.-M. Lee, S. Minwalla, M. Rangamani, and N. Seiberg, *Three point functions of chiral operators in D=4, N=4 SYM at large N*, Adv.Theor.Math.Phys.2(1998)697, hep-th/9806074.
- [4] D. Berenstein, J. M. Maldacena, H. Nastase, *Strings in flat space and pp waves from N=4 super Yang-Mills*, JHEP0204(2002)013, hep-th/0202021.
- [5] S. Dobashi, H. Shimada, and T. Yoneya, *Holographic reformulation of string theory on $AdS_5 \times S^5$ background in the PP-wave limit*, Nucl.Phys.B665(2003)94, hep-th/0209251.
- [6] T. Yoneya, *What is holography in the plane wave limit of $AdS(5)/SYM(4)$ correspondence?*, hep-th/0304183 (expanded from the paper published in the Proceedings, Prog. Theor. Phys. 152 (2003) 108).
- [7] S. Dobashi and T. Yoneya, *Resolving the holography in the plane-wave limit of AdS/CFT correspondence*, hep-th/0406225.
- [8] S. Dobashi and T. Yoneya, *Impurity Non-Preserving 3-Point Correlators of BMN Operators from PP-Wave Holography I : Bosonic Excitations*, hep-th/0409058
- [9] M. Asano, Y. Sekino, and T. Yoneya, *PP wave holography for Dp-brane backgrounds*, Nucl.Phys.B678(2004)197, hep-th/0308024; M. Asano and Y. Sekino, *Large N limit of SYM theories with 16 supercharges from superstrings on Dp-brane backgrounds*, hep-th/0405203; M. Asano, *Stringy effect of the holographic correspondence for Dp-brane backgrounds*, hep-th/0408030.
- [10] D.J. Gross, A. Mikhailov, and R. Roiban, *A calculation of the plane wave string Hamiltonian from N=4 super Yang-Mills theory*, JHEP0305(2003)025, hep-th/0208231
- [11] J. Gomis, S. Moriyama, and J. Park *SYM description of SFT Hamiltonian in a pp wave background*, Nucl.Phys.B659(2003)179, hep-th/0210153
- [12] P. Di Vecchia, J.L. Petersen, M. Petrini, R. Russo, and A. Tanzini, *The three string vertex and the AdS/CFT duality in the pp wave limit*, hep-th/0304025.
- [13] R. Roiban, M. Spradlin, and A. Volovich, *On light cone SFT contact terms in a plane wave* JHEP0310(2003)055, hep-th/0211220
- [14] P. Gutjahr and A. Pankiewicz, *New aspects of the BMN correspondence beyond the planar limit*, hep-th/0407098
- [15] G. Grignani, M. Orselli, B.Ramadanovic, G.W. Semenoff, and D. Young, *Divergence Cancellation and Loop Corrections in String Field Theory on a Plane Wave Background*, hep-th/0508126
- [16] J. Gomis, S. Moriyama, and J. Park *SYM description of pp wave string interactions: singlet sector and arbitrary impurities*, Nucl.Phys.B665(2003)49, hep-th/0301250

- [17] C.S. Chu, V.V. Khoze, and G. Travaglini, *BMN operators with vector impurities, $Z(2)$ symmetry and pp waves*, JHEP0306(2003)050, hep-th/0303107
- [18] G. Georgiou, V.V. Khoze, and G. Travaglini, *New tests of the pp wave correspondence*, JHEP0310(2003)049 hep-th/0306234
- [19] G. Georgiou and G. Travaglini, *Fermion BMN operators, the dilatation operator of $N=4$ SYM, and pp wave string interactions*, JHEP0404(2004)001, hep-th/0403188.
- [20] N. Beisert, C. Kristjansen, J. Plefka, G.W. Semenoff, and M. Staudacher, *BMN correlators and operator mixing in $N=4$ super Yang-Mills theory*, Nucl.Phys.B650(2003)125-161, hep-th/0208178
- [21] N.R. Constable, D.Z. Freedman, M. Headrick, and S.Minwalla, *Operator mixing and the BMN correspondence*, JHEP0210(2002)068, hep-th/0209002
- [22] M. Spradlin and A. Volovich, *Superstring interactions in a pp wave background*, Phys.Rev.D66(2002)086004,, hep-th/0204146.
- [23] M. Spradlin and A. Volovich, *Superstring interactions in a pp wave background 2*, JHEP0301(2003)036, hep-th/0206073.
- [24] A. Pankiewicz, *More comments on superstring interactions in the pp wave background*, JHEP0209(2002)056, hep-th/0208209.
- [25] A. Pankiewicz and B. Stefanski,Jr., *PP wave light cone superstring field theory*, Nucl.Phys.B657(2003)79, hep-th/0210246.
- [26] A. Pankiewicz, *An alternative formulation of light cone string field theory on the plane wave*, JHEP0306(2003)047, hep-th/0304232.
- [27] A. Pankiewicz and B. Stefanski,Jr., *On the uniqueness of plane wave string field theory*, hep-th/0308062.
- [28] C.S. Chu, V.V. Khoze, *Correspondence between the three point BMN correlators and the three string vertex on the pp wave*, JHEP0304(2003)014, hep-th/0301036.
- [29] S. Lee and R. Russo, *Holographic cubic vertex in the PP-wave*, Nucl.Phys.B705(2005)296, hep-th/0409261
- [30] S. Dobashi, Work in progress.
- [31] N.R. Constable, D.Z. Freedman, M.Headrick, S. Minwalla, L. Motl, A. Postnikov, and W. Skiba, *PP wave string interactions from perturbative Yang-Mills theory*, JHEP0207(2002)017, hep-th/0205089.
- [32] A. Parnachev and A.V. Ryzhov, *Strings in the near plane wave background and AdS/CFT*, JHEP0210(2002)066, hep-th/0208010.
- [33] U. Gursoy, *Vector operators in the BMN correspondence*, hep-th/0208041.
- [34] N. Beisert, *BMN operators and superconformal symmetry*, Nucl.Phys.B659(2003)79, hep-th/0211032.

- [35] C. Kristjansen, J. Plefka, G.W. Semenoff, and M. Staudacher, *A new double scaling limit of $N=4$ super Yang-Mills theory and pp wave strings*, Nucl.Phys.B643(2002)3, hep-th/0205033.
- [36] D.Z. Freedman, K. Johnson and J.I. Latoree, *Differential regularization and renormalization: a new method of calculation in quantum field theory*, Nucl.Phys.B371(1992)353.
- [37] Y. H. He, J. H. Schwarz, M. Spradlin, and A. Volovich, *Explicit formulas for Neumann coefficients in the plane wave geometry*, Phys.Rev.D67(2003)086005, hep-th/0211198.

See also,

J. Lucietti, S. Schafer-Nameki, and A. Sinha, *On the plane-wave cubic vertex*, Phys.Rev.D70(2004)026005, hep-th/0402185.